

Metric gluing of Brownian and $\sqrt{8/3}$ -Liouville quantum gravity surfaces

Ewain Gwynne Jason Miller
MIT Cambridge

Abstract

In a recent series of works, Miller and Sheffield constructed a metric on $\sqrt{8/3}$ -Liouville quantum gravity (LQG) under which $\sqrt{8/3}$ -LQG surfaces (e.g., the LQG sphere, wedge, cone, and disk) are isometric to their Brownian surface counterparts (e.g., the Brownian map, half-plane, plane, and disk). We identify the metric gluing of certain collections of independent $\sqrt{8/3}$ -LQG surfaces with boundaries identified together according to LQG length. In particular, we show that if two quantum wedges $\mathcal{W}^-, \mathcal{W}^+$ are conformally welded together according to LQG length to obtain another quantum wedge \mathcal{W} decorated by an independent $\text{SLE}_{8/3}$ -type curve, then the $\sqrt{8/3}$ -LQG metric on \mathcal{W} is equal to the metric gluing of the $\sqrt{8/3}$ -LQG metrics on $\mathcal{W}^-, \mathcal{W}^+$. We prove similar statements for a number of other natural gluing operations for $\sqrt{8/3}$ -LQG surfaces.

Our results imply that the metric gluing of two independent instances of the Brownian half-plane along their positive boundaries is isometric to a certain $\sqrt{8/3}$ -LQG wedge decorated by an independent chordal $\text{SLE}_{8/3}$ curve. If one identifies the two sides of the boundary of a single Brownian half-plane, one obtains a certain $\sqrt{8/3}$ -LQG cone decorated by an independent whole-plane $\text{SLE}_{8/3}$. If one identifies the entire boundaries of the two Brownian half-plane instances, one obtains a different $\sqrt{8/3}$ -LQG cone and the interface between them is a two-sided variant of whole-plane $\text{SLE}_{8/3}$.

Combined with another work of the authors, the present work identifies the scaling limit of self-avoiding walk on random quadrangulations with $\text{SLE}_{8/3}$ on $\sqrt{8/3}$ -LQG.

Contents

1	Introduction	2
1.1	Overview	2
1.2	Background on SLE and LQG	4
1.3	Main results	6
1.4	Remarks on metric gluing	9
1.5	Basic notation	10
1.6	Outline	10
2	Metric space preliminaries	10
2.1	Basic definitions for metrics	10
2.2	The $\sqrt{8/3}$ -LQG metric	11
3	The Brownian disk	14
3.1	Brownian disk definition	14
3.2	Area, length, and distance estimates for the Brownian disk	15
4	Metric gluing	20
4.1	LQG geodesics cannot hit the boundary	20
4.2	Metric gluing of two quantum wedges	22
4.3	Metric gluing in the peanosphere	29

1 Introduction

1.1 Overview

For $\gamma \in (0, 2)$, a *Liouville quantum gravity (LQG) surface* is (formally) a random Riemann surface parameterized by a domain $D \subset \mathbb{C}$ whose Riemannian metric tensor is $e^{\gamma h(z)} dx \otimes dy$, where h is some variant of the Gaussian free field (GFF) on D and $dx \otimes dy$ denotes the Euclidean metric tensor. Liouville quantum gravity surfaces are conjectured to arise as the scaling limits of various random planar map models; the case $\gamma = \sqrt{8/3}$ corresponds to uniformly random planar maps, and other values of γ are obtained by weighting by the partition function of an appropriate statistical mechanics model (see [She16b, KMSW15, GKMW16] for examples of such statistical mechanics models). This has so far been shown to be the case for $\gamma = \sqrt{8/3}$ with respect to the Gromov-Hausdorff topology in the works [Le 13, Mie13] and [MS15a, MS15b, MS15c, MS16a, MS16b] and for all $\gamma \in (0, 2)$ in the so-called peanosphere sense in [She16b, KMSW15, GKMW16] and [DMS14].

Since h is only a distribution (i.e., generalized function) and does not have well-defined pointwise values, this object does not make rigorous sense. However, it was shown by Duplantier and Sheffield [DS11] that one can rigorously define the volume measure associated with an LQG surface. More specifically, there is a measure μ_h on D , called the γ -LQG measure, which is the a.s. limit of regularized versions of $e^{\gamma h(z)} dz$, where dz is the Euclidean volume form. One can similarly define the γ -LQG length measure ν_h on certain curves in D , including ∂D and Schramm's [Sch00] SLE $_{\kappa}$ -type curves for $\kappa = \gamma^2$ [She16a].

The LQG measures μ_h and ν_h are conformally covariant in the following sense. If $D, \tilde{D} \subset \mathbb{C}$, $\phi : \tilde{D} \rightarrow D$ is a conformal map, and we set

$$\tilde{h} = h \circ \phi + Q \log |\phi'|, \quad \text{where} \quad Q = \frac{2}{\gamma} + \frac{\gamma}{2} \quad (1.1)$$

then μ_h is the pushforward of $\mu_{\tilde{h}}$ and ν_h is the pushforward of $\nu_{\tilde{h}}$ under ϕ [DS11, Proposition 2.1]. Hence it is natural to view pairs (D, h) and (\tilde{D}, \tilde{h}) which differ by a transformation as in (1.1) as different parameterizations of the same LQG surface.

In the recent works [MS15a, MS15b, MS15c, MS16a, MS16b], Miller and Sheffield showed that in the special case when $\gamma = \sqrt{8/3}$, an LQG surface is equipped with a natural metric (distance function) \mathfrak{d}_h , so can be interpreted as a random metric space. This metric \mathfrak{d}_h is also invariant under transformations of the form (1.1), so is a functional of the quantum surface itself, not the particular choice of parameterization. We will review the construction of \mathfrak{d}_h in Section 2.2.

In this paper we will be interested in several particular types of $\sqrt{8/3}$ -LQG surfaces which are defined in [DMS14]. These include quantum spheres, which are finite-volume LQG surfaces (i.e., the total mass of the $\sqrt{8/3}$ -LQG measure μ_h is finite) parameterized by the Riemann sphere $\mathbb{C} \cup \{\infty\}$; quantum disks, which are finite-volume LQG surfaces with boundary; α -quantum cones for $\alpha < Q$ (with Q as in (1.1) for $\gamma = \sqrt{8/3}$), which are infinite-volume LQG surfaces homeomorphic to \mathbb{C} ; and α -quantum wedges for $\alpha < Q + \frac{1}{2}\sqrt{8/3}$, which are infinite-volume LQG surfaces with infinite boundary. We will review the definitions of these particular types of quantum surfaces in Section 1.2.2 below.

The *Brownian map* [Le 13, Mie13] is a continuum random metric space which arises as the scaling limit of uniform random planar maps, and which is constructed using a continuum analog of the Schaeffer bijection [Sch97]. We refer to the surveys [Le 14, Mie09] for more details about this object. One can also define Brownian surfaces with other topologies, such as the Brownian half-plane, which is the scaling limit of the uniform infinite half-planar quadrangulations [GM16b, BMR16]; the Brownian plane [CL14], which is the scaling limit of the uniform infinite planar quadrangulation; and the Brownian disk [BM15], which is the scaling limit of finite quadrangulations with boundary. It is shown in [MS16a, Corollary 1.5] (or [GM16b, Proposition 1.10] in the half-plane case) that each of these Brownian surfaces is isometric to a certain special $\sqrt{8/3}$ -LQG surface, i.e. one can couple the two random metric spaces in such a way that there a.s. exists an isometry between them. It is shown in [MS16b] that in fact that the $\sqrt{8/3}$ -LQG structure is a measurable function of the Brownian surface structure and it follows from the construction in [MS15b, MS16a] that the converse holds). In particular,

- The Brownian map is isometric to the quantum sphere;
- The Brownian disk is isometric to the quantum disk;

- The Brownian plane is isometric to the $\sqrt{8/3}$ -quantum cone;
- The Brownian half-plane is isometric to the $\sqrt{8/3}$ -quantum wedge.

Furthermore, the isometries are such that the $\sqrt{8/3}$ -LQG area measure is pushed forward to the natural volume measure on the corresponding Brownian surface and (in the case of the disk or half-plane) the $\sqrt{8/3}$ -LQG boundary length measure is pushed forward to the natural boundary length measure on the Brownian disk or half-plane. That is, the spaces are equivalent as metric measure spaces. Hence the construction of the $\sqrt{8/3}$ -LQG metric in [MS15a, MS15b, MS15c, MS16a, MS16b] can be interpreted as:

- Endowing the Brownian map, half-plane, plane, and disk with a canonical conformal structure and
- Constructing many additional random metric spaces (corresponding to other LQG surfaces) which locally look like Brownian surfaces.

The goal of this paper is to study metric space quotients (also known as metric gluings) of $\sqrt{8/3}$ -LQG surfaces glued along their boundaries according to quantum length. The results described in [DMS14, Section 1.6] (see also [She16a]) show that one can *conformally* glue various types of quantum surfaces along their boundaries according to quantum length to obtain different quantum surfaces. In each case, the interface between the glued surfaces is an SLE_κ -type curve with $\kappa \in \{\gamma^2, 16/\gamma^2\}$ (so $\kappa \in \{8/3, 6\}$ when $\gamma = \sqrt{8/3}$). We will show that in the case when the interface is a simple curve, this conformal gluing is the same as the metric gluing. In other words, the $\sqrt{8/3}$ -LQG metric on the glued surface is the metric quotient of the $\sqrt{8/3}$ -LQG metrics on the surfaces being glued. See Section 1.3 for precise statements. We emphasize that we will be considering metric gluings along rough, fractal curves and in general such gluings are not well-behaved. See Section 1.4 for a discussion of the difficulties involved in proving metric gluing statements of the sort we consider here.




Figure 1: A chart of the different components which serve as input into the proof that self-avoiding walk on random quadrangulations converges to $SLE_{8/3}$ on $\sqrt{8/3}$ -LQG. The present article corresponds to the blue box and implies that the embedding into \mathbb{C} via so-called QLE(8/3, 0) of the metric gluing of a pair of Brownian half-planes to each other along their boundary or a single Brownian half-plane to itself along its boundary is described by $\sqrt{8/3}$ -LQG where the interface is an $SLE_{8/3}$ -type path.

In light of the aforementioned relationship between $\sqrt{8/3}$ -LQG surfaces and Brownian surfaces, our results translate into results for Brownian surfaces. In particular, our results imply that each of the Brownian map

and the Brownian plane are metric space quotients of countably many independent Brownian disks glued together along their boundaries; and that when one metrically glues together two independent Brownian surfaces, resulting surface locally looks like a Brownian surface (even at points along the gluing interface). To our knowledge, it is not known how to derive either of these facts directly from the Brownian surface literature. Hence this work can be viewed as an application of $\sqrt{8/3}$ -LQG to the theory of Brownian surfaces. However, our proofs will also make use of certain facts about Brownian surfaces are not obvious directly from the $\sqrt{8/3}$ -LQG perspective (in particular the estimates for the Brownian disk contained in Section 3.2).

The results of this paper also have applications to scaling limits of random quadrangulations decorated by a self-avoiding walk. Indeed, it is known that gluing together two uniformly random quadrangulations with simple boundary along their boundaries according to boundary length (or gluing two boundary arcs of a single uniformly random quadrangulation with simple boundary to each other according to boundary length) produces a uniformly random pair consisting of a quadrangulation decorated by a self-avoiding walk. See [Bet15, Section 8.2] (which builds on [BBG12, BG09]) for the case of finite quadrangulations with simple boundary and [Car15, Part III], [CC16] for the case of the uniform infinite half-planar quadrangulation with simple boundary (UIHPQ_S). The results of the present work combined with [GM16a, GM16b] imply that the scaling limit of a uniform infinite planar or half-planar quadrangulation decorated by a self-avoiding walk is an appropriate $\sqrt{8/3}$ -LQG surface decorated by an SLE_{8/3}-type curve. See Figure 1 for a schematic diagram of how the different works of the authors fit together with the existing literature to deduce this result.

Acknowledgements E.G. was supported by the U.S. Department of Defense via an NDSEG fellowship. E.G. also thanks the hospitality of the Statistical Laboratory at the University of Cambridge, where this work was started. J.M. thanks Institut Henri Poincaré for support as a holder of the Poincaré chair, during which this work was completed.

1.2 Background on SLE and LQG

Before we state our main results, we give a brief review of some definitions related to SLE and LQG which are needed for the statements.

1.2.1 SLE and space-filling SLE

Let $\kappa > 0$ (here we will only need the case $\kappa = 8/3$) and let $\underline{\rho} = (\rho_1, \dots, \rho_n)$ be a finite vector of weights. Also let $D \subset \mathbb{C}$ be a simply connected domain and let $x, y, z_1, \dots, z_n \in D \cup \partial D$. A *chordal* SLE _{κ} ($\underline{\rho}$) from x to y in D is a variant of chordal SLE _{κ} from x to y in D which has additional *force points* at z_1, \dots, z_n of weights ρ_1, \dots, ρ_n , respectively. It is a conformally invariant process which satisfies the so-called domain Markov process where one has to keep track of the extra marked points. Chordal SLE _{κ} ($\underline{\rho}$) processes were first introduced in [LSW03, Section 8.3]. See also [SW05] and [MS16d, Section 2.2]. In this paper we will primarily be interested in chordal SLE _{κ} ($\rho^L; \rho^R$) with two force points of weight ρ^L and ρ^R located immediately to the left and right of the starting point, respectively. Such a process is well-defined provided $\rho^L, \rho^R > -2$ [MS16d, Section 2.2]. We also recall the definition of whole-plane SLE _{κ} (ρ) for $\rho > -2$ [MS13, Section 2.1].

For $\kappa \in (4, 8)$ (here we will only need the case $\kappa = 6$), *whole-plane space-filling SLE _{κ} from ∞ to ∞* is a variant of SLE _{κ} which travels from ∞ to ∞ in \mathbb{C} and iteratively fills in bubbles as it disconnects them from ∞ (so in particular is not a Loewner evolution). As explained in [DMS14, Footnote 9], whole-plane space-filling SLE _{κ} can be sampled as follows.

1. Let η^L and η^R be the flow lines of a whole-plane GFF started from 0 with angles $\pi/2$ and $-\pi/2$, respectively, in the sense of [MS13]. Note that by [MS13, Theorem 1.1], η^L has the law of a whole-plane SLE _{$16/\kappa$} ($2 - 16/\kappa$) from 0 to ∞ and by [MS13, Theorem 1.11], the conditional law of η^R given η^L is that of a concatenation of independent chordal SLE _{$16/\kappa$} ($-8/\kappa; -8/\kappa$) processes in the connected components of $\mathbb{C} \setminus \eta^L$.
2. Conditional on η^L and η^R , concatenate a collection of independent chordal space-filling SLE _{κ} processes, one in each connected component of $\mathbb{C} \setminus (\eta^L \cup \eta^R)$. Such processes are defined in [MS13, Sections 1.1.3 and 4.3] and can be obtained from ordinary chordal SLE _{κ} by, roughly speaking, iteratively filling in the bubbles it disconnects from ∞ .

1.2.2 Liouville quantum gravity surfaces

In this subsection we will give a brief review of Liouville quantum gravity (LQG) surfaces, as introduced in [DS11, She16a, DMS14]. Such surfaces are defined for all $\gamma \in (0, 2)$, but in this paper we will consider only the case when $\gamma = \sqrt{8/3}$, since this is the only case for which an LQG surface has a rigorously defined metric space structure (which we will review in Section 2.2).

A $\sqrt{8/3}$ -LQG surface is an equivalence class of pairs (D, h) , where $D \subset \mathbb{C}$ is an open set and h is some variant of the Gaussian free field (GFF) [She07, SS13, She16a, MS16d, MS13] on D . Two pairs (D, h) and (\tilde{D}, \tilde{h}) are declared to be equivalent if there is a conformal map $\phi : \tilde{D} \rightarrow D$ such that

$$\tilde{h} = h \circ \phi + Q \log |\phi'|, \quad \text{where} \quad Q = \frac{2}{\gamma} + \frac{\gamma}{2} = \sqrt{\frac{3}{2}} + \sqrt{\frac{2}{3}}. \quad (1.2)$$

The particular choice of distribution h is referred to as the *embedding* of the quantum surface (into D). One can also define quantum surfaces with $k \in \mathbb{N}$ marked points in $D \cup \partial D$ (with points in ∂D viewed as prime ends) by requiring the map ϕ in (1.2) to take the marked points of one surface to the corresponding marked points of the other.

It is shown in [DS11] that a $\sqrt{8/3}$ -LQG surface has a natural area measure μ_h , which is a limit of regularized versions of $e^{\sqrt{8/3}h} dz$, where dz denotes Lebesgue measure on D . Furthermore, there is a natural length measure ν_h which is defined on certain curves in D (including ∂D and SLE $_{8/3}$ -type curves which are independent from h [She16a]), and which is a limit of regularized versions of $e^{\sqrt{8/3}h/2} |dz|$, where $|dz|$ is the Euclidean length element. The measures μ_h and ν_h are invariant under transformations of the form (1.2) (see [DS11, Proposition 2.1] and its length measure analog).

In this paper we will be interested in several specific types of $\sqrt{8/3}$ -LQG surfaces which are defined and studied in [DMS14]. Let Q be as in (1.2). For $\alpha \leq Q$, an α -quantum wedge is a doubly-marked quantum surface $(\mathbb{H}, h, 0, \infty)$ which can be obtained from a free-boundary GFF on \mathbb{H} plus $-\alpha \log |\cdot|$ by “zooming in near 0” so as to fix the additive constant in a canonical way. See [DMS14, Section 4.2] for a precise definition. Quantum wedges in the case when $\alpha \leq Q$ are called *thick wedges* because they describe surfaces homeomorphic to \mathbb{H} . Bounded neighborhoods of the marked point at 0 a.s. have finite mass while neighborhoods of ∞ a.s. have infinite mass.

For $\alpha \in (Q, Q + \sqrt{2/3})$, an α -quantum wedge is an ordered Poissonian collection of doubly-marked quantum surfaces, each with the topology of the disk (the two marked points correspond to the points $\pm\infty$ in the infinite strip in [DMS14, Definition 4.19]). The individual surfaces are called *beads* of the quantum wedge. One can also consider a single bead of an α -quantum wedge conditioned on its quantum area and/or its left and right quantum boundary lengths. See [DMS14, Section 4.4] for more details. Quantum wedges in the case when $\alpha \in (Q, Q + \sqrt{2/3})$ are called *thin wedges* (because they are not homeomorphic to \mathbb{H}). It is sometimes convenient to parameterize the set of quantum wedges by a different parameter \mathfrak{w} , called the *weight*, which is given by

$$\mathfrak{w} = \sqrt{\frac{8}{3}} \left(\sqrt{\frac{2}{3}} + Q - \alpha \right) > 0. \quad (1.3)$$

The reason why the weight parameter is convenient is that it is additive under the gluing and cutting operations for quantum wedges and quantum cones studied in [DMS14].

For $\alpha < Q$, an α -quantum cone, defined in [DMS14, Definition 4.13], is a doubly-marked quantum surface $(\mathbb{C}, h, 0, \infty)$ obtained from a whole-plane GFF plus $-\alpha \log |\cdot|$ by “zooming in near 0” (so as to fix the additive constant in a canonical way) in a similar manner to how a thick wedge is obtained from a free-boundary GFF with a log singularity. The *weight* of an α -quantum cone is given by

$$\mathfrak{w} = 2\sqrt{\frac{8}{3}} (Q - \alpha). \quad (1.4)$$

A *quantum sphere* is a finite-volume quantum surface (\mathbb{C}, h) defined in [DMS14, Definition 4.21], which is often taken to have fixed quantum area. One can also consider quantum spheres with one, two, or three marked points, which we take to be sampled uniformly and independently from the $\sqrt{8/3}$ -LQG area

measure μ_h . Note that the marked points in [DMS14, Definition 4.21] correspond to the points $\pm\infty$ in the cylinder, and are shown to be sampled uniformly and independently from μ_h in [DMS14, Proposition 5.15].

A *quantum disk* is a finite-volume quantum surface with boundary (\mathbb{D}, h) defined in [DMS14, Definition 4.21], which can be taken to have fixed area, boundary length, or both. A *singly (resp. doubly) marked quantum disk* is a quantum disk together with one (resp. two) marked points in $\partial\mathbb{D}$ sampled uniformly (and independently) from the $\sqrt{8/3}$ -LQG boundary length measure ν_h . Note that the marked points in [DMS14, Definition 4.21] correspond to the points $\pm\infty$ in the infinite strip, and are shown to be sampled uniformly and independently from μ_h in [DMS14, Proposition 5.7]. It follows from the definitions in [DMS14, Section 4.4] that a doubly-marked quantum disk has the same law as a single bead of a (weight-2/3) $\sqrt{6}$ -quantum wedge if we condition on quantum area and left/right quantum boundary lengths (note that this is only true for $\gamma = \sqrt{8/3}$).

Suppose $\mathfrak{w}^-, \mathfrak{w}^+ > 0$ and $\mathfrak{w} = \mathfrak{w}^- + \mathfrak{w}^+$. It is shown in [DMS14, Theorem 1.9] that if one cuts a weight- \mathfrak{w} quantum wedge by an independent chordal $\text{SLE}_{8/3}(\mathfrak{w}^- - 2; \mathfrak{w}^+ - 2)$ curve (or a concatenation of such curves in the thin wedge case) then one obtains a weight- \mathfrak{w}^- quantum wedge and an independent- \mathfrak{w}^+ quantum wedge which can be glued together according to quantum boundary length to recover the original weight- \mathfrak{w} quantum wedge. Similarly, by [DMS14, Theorem 1.12], if one cuts a weight- \mathfrak{w} quantum cone by an independent whole-plane $\text{SLE}_{8/3}(\mathfrak{w} - 2)$ curve, then one obtains a weight- \mathfrak{w} quantum wedge whose left and right boundaries can be glued together according to quantum length to recover the original weight- \mathfrak{w} quantum cone.

1.3 Main results

In this section we will state the main results of this paper. Our first two theorems state that whenever we have a conformal welding of two or more $\sqrt{8/3}$ -LQG surfaces along a simple $\text{SLE}_{8/3}$ -type curve as in [She16a, DMS14], the metric on the welding of the surfaces is equal to the metric space quotient of the surfaces being welded together according to boundary length. We start by addressing the case of two quantum wedges glued together to obtain another quantum wedge.

Theorem 1.1. *Let $\mathfrak{w}^-, \mathfrak{w}^+ > 0$ and let $\mathfrak{w} := \mathfrak{w}^- + \mathfrak{w}^+$. If $\mathfrak{w} \geq 4/3$, let $(\mathbb{H}, h, 0, \infty)$ be a weight- \mathfrak{w} quantum wedge (recall (1.3)). If $\mathfrak{w} \in (0, 4/3)$, let $(\mathbb{H}, h, 0, \infty)$ be a single bead of a weight- \mathfrak{w} quantum wedge with area $\mathfrak{a} > 0$ and left/right boundary lengths $\mathfrak{l}^-, \mathfrak{l}^+ > 0$. Let η be an independent chordal $\text{SLE}_{8/3}(\mathfrak{w}^- - 2; \mathfrak{w}^+ - 2)$ from 0 to ∞ in \mathbb{H} . Let W^- (resp. W^+) be the region lying to the left (resp. right) of η and let \mathcal{W}^- (resp. \mathcal{W}^+) be the quantum surface obtained by restricting h to W^- (resp. W^+). Let \mathcal{U}^- (resp. \mathcal{U}^+) be the ordered sequence of connected components of the interior of W^- (resp. W^+). Let \mathfrak{d}_h be the $\sqrt{8/3}$ -LQG metric induced by h . For $U \in \mathcal{U}^\pm$ let $\mathfrak{d}_{h|_U}$ be the $\sqrt{8/3}$ -LQG metric induced by $h|_U$, extended by continuity to ∂U . Almost surely, the metric space $(\mathbb{H}, \mathfrak{d}_h)$ is the quotient of the disjoint union of the metric spaces $(\overline{U}, \mathfrak{d}_{h|_U})$ for $U \in \mathcal{U}^- \cup \mathcal{U}^+$ under the natural identification (i.e., according to quantum length).*

See Figure 2 (resp. Figure 3) for an illustration of the statement of Theorem 1.1 in the case when $\mathfrak{w}^-, \mathfrak{w}^+ \geq 4/3$ (resp. $\mathfrak{w}^-, \mathfrak{w}^+ < 4/3$). The reason why the metrics $\mathfrak{d}_{h|_U}$ for $U \in \mathcal{U}^- \cup \mathcal{U}^+$ extend continuously to ∂U is explained in Lemma 2.5 below.

In the setting of Theorem 1.1, [DMS14, Theorem 1.9] implies that the quantum surfaces \mathcal{W}^- and \mathcal{W}^+ are independent, the former is a weight- \mathfrak{w}^- quantum wedge (or a collection of beads of such a wedge if $\mathfrak{w}^- < 4/3$), and the latter is a weight- \mathfrak{w}^+ quantum wedge (or a collection of beads of such a wedge if $\mathfrak{w}^+ < 4/3$). Note in particular that $\mathcal{U}^- = W^-$ if $\mathfrak{w}^- \geq 4/3$ and \mathcal{U}^- is countably infinite if $\mathfrak{w}^- < 4/3$, and similarly for \mathcal{U}^+ .

The wedges (or subsets of wedges) \mathcal{W}^- and \mathcal{W}^+ are glued together according to $\sqrt{8/3}$ -LQG boundary length [She16a, Theorem 1.3]. Hence Theorem 1.1 says that one can metrically glue a (subset of a) weight- \mathfrak{w}^- quantum wedge and a (subset of a) weight- \mathfrak{w}^+ quantum wedge together according to quantum boundary length to get a (bead of a) weight- \mathfrak{w} quantum wedge.

The quotient metric space in Theorem 1.1 is an LQG surface so locally looks the same as a Brownian surface. In fact, the quotient metric \mathfrak{d}_h is independent from the gluing interface η . Hence if one takes a metric quotient of two quantum wedges, it is not possible to recover the two original wedges from the quotient space.

We note that by local absolute continuity, Theorem 1.1 implies that a metric space quotient of any pair of independent $\sqrt{8/3}$ (equivalently Brownian) surfaces with boundary glued together according to boundary length also locally looks like a Brownian surface. This applies in particular if one metrically glues together




Figure 2: **Left:** Illustration of the statement of Theorem 1.1 in the case when $\mathfrak{w}^-, \mathfrak{w}^+ \geq 4/3$ (which is the setting of Corollary 1.2). The metric space $(\mathbb{H}, \mathfrak{d}_h)$, which is a weight- \mathfrak{w} quantum wedge, is the metric quotient of the disjoint union of the independent weight- \mathfrak{w}^\pm quantum wedges $(W^\pm, \mathfrak{d}_{h|_{W^\pm}})$ glued together along half of their boundaries according to quantum length. The gluing interface η is a chordal $\text{SLE}_{8/3}(\mathfrak{w}^- - 2; \mathfrak{w}^+ - 2)$ curve. In the case that $\mathfrak{w}^-, \mathfrak{w}^+ = 2$ so that $(W^\pm, \mathfrak{d}_{h|_{W^\pm}})$ are Brownian half-plane instances, η is exactly a chordal $\text{SLE}_{8/3}$. **Middle:** Illustration of the statement of Theorem 1.3. The metric space $(\mathbb{H}, \mathfrak{d}_h)$, which is a weight- \mathfrak{w} quantum cone, is the quotient of the metric space $(U, \mathfrak{d}_{h|_U})$, which is a weight- \mathfrak{w} quantum wedge, under the quotient map which identifies the two sides of the boundary of U according to quantum length. The gluing interface η is a whole-plane $\text{SLE}_{8/3}(\mathfrak{w} - 2)$ curve. In the case that $\mathfrak{w} = 2$ so that $(\mathbb{H}, \mathfrak{d}_h)$ is a Brownian half-plane instance, η is exactly a whole-plane $\text{SLE}_{8/3}$. **Right:** Illustration of Corollary 1.5. Two weight-2 quantum wedges (Brownian half-planes) are glued together along their whole boundaries according to quantum length to obtain a weight-4 quantum cone. The gluing interface is a pair of non-intersecting $\text{SLE}_{8/3}$ -type curves.

two Brownian disks according to boundary length. We emphasize that it is not at all clear from the definition of a Brownian disk that gluing two such objects together produces something which locally looks like a Brownian disk near the gluing interface.

We will now state a corollary of Theorem 1.1 which identifies the law of two independent Brownian half-plane instances glued together according to boundary length on their positive boundaries. This result will allow us interpret the main result of [GM16a] as a convergence result for self-avoiding walk on random quadrangulations to $\text{SLE}_{8/3}$ on $\sqrt{8/3}$ -LQG.

Corollary 1.2. *Suppose that we have two independent instances of the Brownian half-plane. Then the metric quotient obtained by gluing the two surfaces together according to boundary length on their positive boundaries is isometric to the $\sqrt{8/3}$ -LQG metric associated with a weight-4 quantum wedge. Moreover, the interface between the two Brownian half-plane instances is a chordal $\text{SLE}_{8/3}$ curve.*

Corollary 1.2 is immediate from Theorem 1.1 since the Brownian half-plane is isometric to the $\sqrt{8/3}$ (weight 2) quantum wedge.

Our next theorem concerns a quantum wedge glued to itself to obtain a quantum cone and is illustrated in the middle of Figure 2.

Theorem 1.3. *Let $\mathfrak{w} \geq 4/3$, let $(\mathbb{C}, h, 0, \infty)$ be a weight- \mathfrak{w} quantum cone (recall (1.4)), and let \mathfrak{d}_h be the $\sqrt{8/3}$ -LQG measure induced by h . Let η be a whole-plane $\text{SLE}_{8/3}(\mathfrak{w} - 2)$ process from 0 to ∞ . Let $U = \mathbb{C} \setminus \eta$ and let $\mathfrak{d}_{h|_U}$ be the $\sqrt{8/3}$ -LQG measure induced by $h|_U$, extended by continuity to ∂U (viewed as a collection of prime ends). Then a.s. $(\mathbb{C}, \mathfrak{d}_h)$ is the metric quotient of $(U, \mathfrak{d}_{h|_U})$ under the natural identification of the two sides of η .*

Note that [DMS14, Theorem 1.12] implies that the surface $(U, h|_U, 0, \infty)$ has the law of a weight- \mathfrak{w} quantum wedge. Furthermore, the boundary arcs of this quantum wedge lying to the left and right of 0 are




Figure 3: **Left:** Illustration of the statement of Theorem 1.1 in the case when $\mathfrak{w} \geq 4/3$ and $\mathfrak{w}^-, \mathfrak{w}^+ < 4/3$. The metric space $(\mathbb{H}, \mathfrak{d}_h)$, which is a weight- \mathfrak{w} quantum wedge, is the metric quotient of the disjoint union of the metric spaces $(U, \mathfrak{d}_{h|_U})$ for $U \in \mathcal{U}^- \cup \mathcal{U}^+$, which are the beads of a pair of independent quantum wedges of weights \mathfrak{w}^- and \mathfrak{w}^+ , glued together according to quantum boundary length. The gluing interface is a chordal $\text{SLE}_{8/3}(\mathfrak{w}_1 - 2; \mathfrak{w}_2 - 2)$ curve η from 0 to ∞ in \mathbb{H} , which a.s. hits both $(-\infty, 0]$ and $[0, \infty)$. **Right:** Illustration of the statement of Theorem 1.6. The boundary of a space-filling SLE_6 curve stopped when it hits 0 divides \mathbb{C} into two independent weight-2/3 quantum wedges (green and blue), whose beads are independent quantum disks (Brownian disks) when we condition on boundary length. The weight 4/3 quantum cone metric space $(\mathbb{C}, \mathfrak{d}_h)$, which is a Brownian plane, is the metric quotient of these Brownian disks glued together according to boundary length.

glued together according to $\sqrt{8/3}$ -LQG boundary length [She16a, Theorem 1.3]. Hence Theorem 1.3 implies that one can metrically glue the left and right boundaries of a weight- \mathfrak{w} quantum wedge together according to quantum length to get a weight- \mathfrak{w} quantum cone. Similar absolute continuity remarks to the ones made after the statement of Theorem 1.1 apply in the setting of Theorem 1.3.

Requiring that $\mathfrak{w} \geq 4/3$ in Theorem 1.3 is equivalent to requiring that this wedge is thick, equivalently the curve η is simple and the set U is connected. We expect that one can also metrically glue the beads of a weight- \mathfrak{w} quantum wedge for $\mathfrak{w} \in (0, 4/3)$ together according to quantum length to get a weight- \mathfrak{w} quantum cone, but will not treat this case in the present work.

We next state an analog of Corollary 1.2 in the setting of Theorem 1.3, which again follows from the fact that the weight 2 quantum wedge is the same as the Brownian half-plane.

Corollary 1.4. *The metric space obtained by gluing the positive and negative boundaries of an instance of the Brownian half-plane together according to boundary length is isometric to the $\sqrt{8/3}$ -LQG metric associated with a weight-2 quantum cone and the interface is a whole-plane $\text{SLE}_{8/3}$ curve.*

Iteratively applying Theorems 1.1 and 1.3 can give us metric gluing statements for when we cut a quantum surface by multiple $\text{SLE}_{8/3}$ -type curves. One basic application of this fact is the following variant of Corollary 1.2 in which one identifies the entire boundary of the two Brownian half-plane instances together.

Corollary 1.5. *Suppose that we have two independent instances of the Brownian half-plane. Then the metric quotient obtained by gluing the two surfaces together according to boundary length is isometric to the $\sqrt{8/3}$ -LQG metric associated with a weight-4 quantum cone. Moreover, the interface between the two*

Brownian half-plane instances is an $SLE_{8/3}$ -type curve which can be sampled as follows. First sample a whole-plane $SLE_{8/3}(2)$ curve η_1 from 0 to ∞ ; then, conditional on η_1 , sample a chordal $SLE_{8/3}$ curve η_2 from 0 to ∞ in $\mathbb{C} \setminus \eta_1$.

We remark that the interface in Corollary 1.5 can also be described by a pair of GFF flow lines [MS16d, MS13].

Corollary 1.5 is obtained by applying Corollary 1.2 to glue the positive boundaries of the two Brownian half-planes and then applying Theorem 1.3 to glue the two sides of the resulting weight-4 quantum wedge.

Another result which can be obtained from multiple applications of Theorems 1.1 and 1.3 (together with the description of the law of whole-plane space-filling SLE_6 in Section 1.2.1) is the following statement, which applies in the setting of the peanosphere construction [DMS14, Theorem 1.13].

Theorem 1.6. *Let $(\mathbb{C}, h, 0, \infty)$ be a $\sqrt{8/3}$ -quantum cone (weight-4/3) and let η' be an independent whole-plane space-filling SLE_6 parameterized by quantum mass with respect to h and satisfying $\eta'(0) = 0$. Let \mathcal{U}^- (resp. \mathcal{U}^+) be the set of connected components of the interior of $\eta'((-\infty, 0])$ (resp. $\eta'([0, \infty))$). Let \mathfrak{d}_h be the $\sqrt{8/3}$ -LQG measure induced by h and for $U \in \mathcal{U}^- \cup \mathcal{U}^+$ let $\mathfrak{d}_{h|_U}$ be the $\sqrt{8/3}$ -LQG metric induced by $h|_U$, extended by continuity to ∂U . Then a.s. $(\mathbb{C}, \mathfrak{d}_h)$ is the metric quotient of the disjoint union of the metric spaces $(\overline{U}, \mathfrak{d}_{h|_U})$ for $U \in \mathcal{U}^- \cup \mathcal{U}^+$ under the natural identification.*

By [MS16a, Corollary 1.5], the metric space $(\mathbb{C}, \mathfrak{d}_h)$ in Theorem 1.6 is isometric to the Brownian plane. Furthermore, by [DMS14, Theorems 1.9 and 1.12], each of the surfaces $(U, h|_U)$ has the law of a single bead of a weight-2/3 quantum wedge, which has the same law as a quantum disk. Hence each of the metric spaces $(U, \mathfrak{d}_{h|_U})$ is isometric to a Brownian disk. Therefore Theorem 1.6 expresses the Brownian plane as a metric space quotient of a countable collection of Brownian disks glued together according to boundary length.

The following is an analog of Theorem 1.6 in the setting of the finite-volume peanosphere construction [MS15c, Theorem 1.1].

Theorem 1.7. *Let (\mathbb{C}, h, ∞) be a singly marked unit area $\sqrt{8/3}$ -quantum sphere and let η' be an independent whole-plane space-filling SLE_6 parameterized by quantum mass with respect to h . Let $\mathfrak{t} \in [0, 1]$ be sampled uniformly from Lebesgue measure, independent from everything else. Let \mathcal{U}^- (resp. \mathcal{U}^+) be the set of connected components of the interior of $\eta'([0, \mathfrak{t}])$ (resp. $\eta'([\mathfrak{t}, 1])$). Let \mathfrak{d}_h be the $\sqrt{8/3}$ -LQG measure induced by h and for $U \in \mathcal{U}^- \cup \mathcal{U}^+$ let $\mathfrak{d}_{h|_U}$ be the $\sqrt{8/3}$ -LQG measure induced by $h|_U$, extended by continuity to ∂U . Then a.s. $(\mathbb{C}, \mathfrak{d}_h)$ is the metric quotient of the disjoint union of the metric spaces $(\overline{U}, \mathfrak{d}_{h|_U})$ for $U \in \mathcal{U}^- \cup \mathcal{U}^+$ under the natural identification.*

By [MS16a, Corollary 1.5], the metric space $(\mathbb{C}, \mathfrak{d}_h)$ in Theorem 1.7 is isometric to the Brownian map. Furthermore, by [MS15c, Theorem 7.1] each of the surfaces $(U, h|_U)$ are conditionally independent given their quantum boundary lengths and areas, and each has the law of a single bead of a weight-2/3 quantum wedge (which has the same law as a quantum disk with given boundary length and area) under this conditioning. Hence each of the metric spaces $(U, \mathfrak{d}_{h|_U})$ is isometric to a Brownian disk. Therefore Theorem 1.6 expresses the Brownian map as a metric space quotient of a countable collection of Brownian disks glued together according to boundary length.

1.4 Remarks on metric gluing

Here we will discuss the metric gluing problem which we study in this paper at an abstract level and explain some of the difficulties involved.

We know a priori (see Lemma 2.2 below) that in the setting of either Theorem 1.1 or Theorem 1.3, it holds for each $U \in \mathcal{U}^- \cup \mathcal{U}^+$ that the restriction of the metric $\mathfrak{d}_{h|_U}$ to the open set U coincides with the internal metric of \mathfrak{d}_h on U (whose definition we recall in Section 2.1). However, this fact together with the fact that the gluing interface η is a continuous simple curve is far from implying the statements of Theorems 1.1 and 1.3.

The problem of proving that a metric space cut by a simple curve is the quotient of the internal metrics on the two sides of the curve is similar in spirit to the problem of proving that a curve in \mathbb{C} is *conformally removable* [JS00], which means that any homeomorphism of \mathbb{C} which is conformal off of the curve is in fact conformal on the whole plane. Indeed, proving each involves estimating how much the length of a path (the

image of a straight line in the case of removability or a geodesic in the case of metric gluing) is affected by its crossings of the curve. Moreover, in the setting of LQG and SLE, both the question of removability and the metric gluing problem addressed in this paper are ways to show that the surfaces formed by cutting along an SLE curve together determine the overall surface (see [She16a, DMS14] for further discussion of this point in the case of removability), although we are not aware of a direct relationship between the two concepts.

SLE $_{\kappa}$ -type curves for $\kappa < 4$ are conformally removable since they are boundaries of Hölder domains [RS05, JS00]. However, there is no such simple criterion for metric gluing. Indeed, it is not even enough to know that the gluing interface is Lipschitz, as the following example demonstrates.

Let $X = [0, 1] \times [-1, 1]$ equipped with the Euclidean metric d and let (\tilde{X}, \tilde{d}) be the metric quotient of $X \sqcup [0, 1/2]$ under the map $[0, 1/2] \ni t \mapsto (2t, 0) \in X$. The space (X, d) is the quotient of the $[0, 1] \times [-1, 0]$ and $[0, 1] \times [0, 1]$ equipped with their d -internal metrics (which both coincide with the Euclidean metric) under the natural identification. Furthermore, (X, d) and (\tilde{X}, \tilde{d}) are homeomorphic (via the obvious identification) and the internal metrics of d and \tilde{d} on the two sides $[0, 1] \times [-1, 0)$ and $[0, 1] \times (0, 1]$ of $[0, 1] \times \{0\}$ coincide. However, $d \neq \tilde{d}$ since points near the interface $[0, 1] \times \{0\}$ are almost twice as far apart with respect to d as with respect to \tilde{d} .

In the example above, \tilde{d} -geodesics between points near $[0, 1] \times \{0\}$ spend most of their time in $[0, 1] \times \{0\}$. A major part of the proof of Theorems 1.1 and 1.3 will consist of controlling how often a \mathfrak{d}_h geodesic hits η and how it behaves near η , in order to rule out the aforementioned type of behavior. However, since the gluing interface η in our setting is only Hölder continuous (with respect to the one-sided and two-sided metrics), not Lipschitz like the curve in the above example, additional estimates are needed as well.

1.5 Basic notation

Here we record some basic notation which we will use throughout this paper.

Notation 1.8. For $a < b \in \mathbb{R}$ and $c > 0$, we define the discrete intervals $[a, b]_{c\mathbb{Z}} := [a, b] \cap (c\mathbb{Z})$ and $(a, b)_{c\mathbb{Z}} := (a, b) \cap (c\mathbb{Z})$.

Notation 1.9. If a and b are two quantities, we write $a \preceq b$ (resp. $a \succeq b$) if there is a constant C (independent of the parameters of interest) such that $a \leq Cb$ (resp. $a \geq Cb$). We write $a \asymp b$ if $a \preceq b$ and $a \succeq b$.

Notation 1.10. If a and b are two quantities which depend on a parameter x , we write $a = o_x(b)$ (resp. $a = O_x(b)$) if $a/b \rightarrow 0$ (resp. a/b remains bounded) as $x \rightarrow 0$ (or as $x \rightarrow \infty$, depending on context).

Unless otherwise stated, all implicit constants in \asymp , \preceq , and \succeq and $O_x(\cdot)$ and $o_x(\cdot)$ errors involved in the proof of a result are required to depend only on the auxiliary parameters that the implicit constants in the statement of the result are allowed to depend on.

1.6 Outline

The remainder of this article is structured as follows. In Section 2, we review the definitions of internal metrics and quotient metrics (i.e., metric gluings) as well as the construction of the $\sqrt{8/3}$ -LQG metric from [MS15b, MS16a, MS16b]. In Section 3, we recall the definition of the Brownian disk from [BM15] and prove some estimates for this metric space which will be needed for the proofs of our main results. In Section 4 we prove our main results using the estimates from Section 3 and some facts about SLE and LQG.

2 Metric space preliminaries

2.1 Basic definitions for metrics

In this paper we will consider a variety of metric spaces, including subsets of \mathbb{C} equipped with the Euclidean metric, $\sqrt{8/3}$ -LQG surfaces equipped with the metric induced by QLE(8/3, 0), and various Brownian surfaces equipped with their intrinsic metric. Here we introduce some notation to distinguish these metric spaces and recall some basic constructions for metric spaces.

Definition 2.1. Let (X, d_X) be a metric space. For $A \subset X$ we write $\text{diam}(A; d_X)$ for the supremum of the d_X -distance between pairs of points in A . For $r > 0$, we write $B_r(A; d_X)$ for the set of $x \in X$ with $d_X(x, A) < r$. If $A = \{y\}$ is a singleton, we write $B_r(\{y\}; d_X) = B_r(y; d_X)$. For $A \subset \mathbb{C}$, we write $B_r(A)$ for the set of points lying at Euclidean distance (strictly) less than r from A .

Recall that a *pseudometric* on a set X is a function $d_X : X \times X \rightarrow [0, \infty)$ which satisfies all of the conditions in the definition of a metric except that possibly $d_X(x, y) = 0$ for $x \neq y$.

Let (X, d_X) be a space equipped with a pseudometric d_X , let \sim be an equivalence relation on X , and let $\overline{X} = X/\sim$ be the corresponding topological quotient space. For equivalence classes $\overline{x}, \overline{y} \in \overline{X}$, let $\mathcal{Q}(\overline{x}, \overline{y})$ be the set of finite sequences $(x_1, y_1, \dots, x_n, y_n)$ of elements of X such that $x_1 \in \overline{x}$, $y_n \in \overline{y}$, and $y_i \sim x_{i+1}$ for each $i \in [1, n-1]_{\mathbb{Z}}$. Let

$$\overline{d}_X(\overline{x}, \overline{y}) := \inf_{\mathcal{Q}(\overline{x}, \overline{y})} \sum_{i=1}^n d_X(x_i, y_i). \quad (2.1)$$

Then \overline{d}_X is a pseudometric on \overline{X} , which we call the *quotient pseudometric*.

For a curve $\gamma : [a, b] \rightarrow X$, the d_X -length of γ is defined by

$$\text{len}(\gamma; d_X) := \sup_P \sum_{i=1}^{\#P} d_X(\gamma(t_i), \gamma(t_{i-1}))$$

where the supremum is over all partitions $P : a = t_0 < \dots < t_{\#P} = b$ of $[a, b]$. Note that the d_X -length of a curve may be infinite.

Suppose $Y \subset X$. The *internal metric* of d_X on Y is defined by

$$d_Y(x, y) := \inf_{\gamma \subset Y} \text{len}(\gamma; d_X), \quad \forall x, y \in Y \quad (2.2)$$

where the infimum is over all curves in Y from x to y . The function d_Y satisfies all of the properties of a metric on Y except that it may take infinite values.

2.2 The $\sqrt{8/3}$ -LQG metric

Suppose (D, h) is a $\sqrt{8/3}$ -LQG surface. In [MS15b, MS16a, MS16b], it is shown that if h is some variant of the GFF on D , then h induces a metric \mathfrak{d}_h on D (which in many cases extends to a metric on $D \cup \partial D$). The construction of this metric builds on the results of [DMS14, MS15a, MS15c, MS16c].

In the special case when (D, h) is a quantum sphere, the metric space (D, \mathfrak{d}_h) is isometric to the Brownian map [MS16a, Theorem 1.4]. It is shown in [MS16a, Corollary 1.5] that the metric space (D, \mathfrak{d}_h) is isometric to a Brownian surface in two additional cases: when (D, h) is a quantum disk we obtain a Brownian disk [BM15] and when (D, h) is a $\sqrt{8/3}$ -quantum cone we obtain a Brownian plane [CL14]. It is shown in [GM16b] that the Brownian half-plane is isometric to the $\sqrt{8/3}$ -quantum wedge. Hence $\sqrt{8/3}$ -LQG surfaces can be viewed as Brownian surfaces equipped with a conformal structure.

For the convenience of the reader, we give in this section a review of the construction of the metric \mathfrak{d}_h and note some basic properties which it satisfies.

2.2.1 The metric on a quantum sphere

The $\sqrt{8/3}$ -LQG metric is first constructed in the case of a $\sqrt{8/3}$ -LQG sphere (\mathbb{C}, h) . Conditional on h , let \mathcal{C} be a countable collection of i.i.d. points sampled uniformly from the $\sqrt{8/3}$ -LQG area measure μ_h . One first defines for $z, w \in \mathcal{C}$ a QLE(8/3, 0) growth process $\{\Gamma_t^{z,w}\}_{t \geq 0}$ started from z and targeted at w , which is a continuum analog of first passage percolation on a random planar map [MS16c, Section 2.2].¹ This is accomplished as follows. Let $\delta > 0$ and let η_0^δ be a whole-plane SLE₆ from z to w sampled independently

¹It is expected that the process $\Gamma_t^{z,w}$ is a re-parameterization of a whole-plane version of the QLE(8/3, 0) processes considered in [MS16c], which are parameterized by capacity instead of by quantum natural time. However, this has not yet been proven to be the case.

from h and then run for δ units of quantum natural time as determined by h [DMS14, Definition 7.2]. For $t \in [0, \delta]$, let $\Gamma_t^{z,w,\delta} := \eta_0^\delta([0, \tau_0^\delta \wedge \delta])$, where τ is the first time η_0^δ hits w .

Inductively, suppose $k \in \mathbb{N}$ and $\Gamma_t^{z,w,\delta}$ has been defined for $t \in [0, k\delta]$. If $w \in \Gamma_{k\delta}^{z,w,\delta}$, let $\Gamma_t^{z,w,\delta} = \Gamma_{k\delta}^{z,w,\delta}$ for each $t \in [k\delta, (k+1)\delta]$. Otherwise, let x_k^δ be sampled uniformly from the $\sqrt{8/3}$ -LQG length measure ν_h restricted to the boundary of the connected component of $\mathbb{C} \setminus \Gamma_t^{z,w,\delta}$ containing w . Let η_k^δ be a radial SLE₆ from x_k^δ to w sampled conditionally independently of h given x_k^δ and $\{\Gamma_s^{z,w,\delta}\}_{s \leq t}$ and parameterized by quantum natural time as determined by h . For $t \in [k\delta, (k+1)\delta]$, let $\Gamma_t^{z,w,\delta} := \eta_k^\delta([0, \tau \wedge (t - k\delta)]) \cup \Gamma_{k\delta}^{z,w,\delta}$, where τ_k^δ is the first time that η_k^δ hits w .

The above procedure defines for each $\delta > 0$ a growing family of sets $\{\Gamma_t^{z,w,\delta}\}_{t \geq 0}$ started from z and stopped when it hits w . It is shown in [MS15b] that (along an appropriately chosen subsequence), one can take an a.s. limit (in an appropriate topology) as $\delta \rightarrow 0$ to obtain a growing family of sets $\{\Gamma_t^{z,w}\}_{t \geq 0}$ from z to w , which we call QLE(8/3,0). It is shown in [MS16a] that the limiting process $\{\Gamma_t^{z,w}\}_{t \geq 0}$ is a.s. determined by h , even though the approximations are not and that the limit does not depend on the choice of subsequence.

For $t \geq 0$, let $X_t^{z,w}$ be the ν_h -length of the boundary of the connected component of $\mathbb{C} \setminus \Gamma_t^{z,w}$ containing w . Let $\sigma_s^{z,w}$ for $r \geq 0$ be defined by

$$r = \int_0^{\sigma_r^{z,w}} \frac{1}{X_t^{z,w}} dt. \quad (2.3)$$

Set $\tilde{\Gamma}_r^{z,w} := \Gamma_{\sigma_r^{z,w}}^{z,w}$. The $\sqrt{8/3}$ -LQG distance between z and w is defined by

$$\mathfrak{d}_h(z, w) := \inf \left\{ r \geq 0 : w \in \tilde{\Gamma}_r^{z,w} \right\}.$$

The time-change (2.3) is natural from the perspective of first passage percolation. Indeed, quantum natural time is the continuum analog of parameterizing a percolation growth by the number of edges traversed, hence the time change (2.3) is the continuum analog of adding edges to the cluster at a rate proportional to boundary length. It is shown in [MS15b] that this function defines a metric on the set \mathcal{C} (which is a.s. dense in \mathbb{C}). It is shown in [MS16a] that \mathfrak{d}_h in fact extends continuously to a metric on all of \mathbb{C} , which is mutually Hölder continuous with respect to the metric on \mathbb{C} induced by the stereographic projection of the standard metric on the Euclidean sphere \mathbb{S}^2 and isometric to the Brownian map. In particular, \mathfrak{d}_h is a geodesic metric.

The re-parameterized QLE(8/3,0) processes $\{\tilde{\Gamma}_r^{z,w}\}_{r \geq 0}$ for $z, w \in \mathcal{C}$ are related to metric balls for \mathfrak{d}_h as follows. For $z, w \in \mathcal{C}$ and $r \geq 0$, the connected component of $\mathbb{C} \setminus \tilde{\Gamma}_r^{z,w}$ containing w is the same as the connected component of $\mathbb{C} \setminus \bar{B}_r(z; \mathfrak{d}_h)$ containing w .

Each of the re-parameterized QLE(8/3,0) hulls $\tilde{\Gamma}_r^{z,w}$ is a local set for h in the sense of [SS13, Section 3.3], and furthermore are determined locally by h . Hence the definition of the metric \mathfrak{d}_h implies that if $U \subset \mathbb{C}$ is a deterministic connected open set, then the quantities $\{\mathfrak{d}_h(z, w) \wedge \mathfrak{d}_h(z, \partial U) : z, w \in U\}$ are a.s. determined by $h|_U$. In particular, the internal metric of \mathfrak{d}_h on U (Section 2.1) is a.s. determined by $h|_U$.

The above metric construction also works with $h + R$ in place of h for any $R \in \mathbb{R}$, in which case [MS16a, Lemma 2.2] yields a scaling property for the metric \mathfrak{d}_h :

$$\mathfrak{d}_{h+R}(z, w) = e^{\sqrt{8/3}R/4} \mathfrak{d}_h(z, w).$$

It is shown in [MS16b] that the $\sqrt{8/3}$ -LQG surface associated with a given Brownian surface is almost surely determined by the metric measure space structure associated with the Brownian surface. This in particular implies that if one is given an instance of the Brownian map, disk, half-plane, or plane, respectively, then there is a measurable way to embed the surface to obtain an instance of a $\sqrt{8/3}$ -LQG sphere, disk, wedge, or cone, respectively. As mentioned above, the construction of the $\sqrt{8/3}$ -LQG metric also implies that the Brownian map, disk, half-plane, or plane structure is a measurable function of the corresponding $\sqrt{8/3}$ -LQG structure. In this way, Brownian and $\sqrt{8/3}$ -LQG surfaces are one and the same.

2.2.2 Metrics on general $\sqrt{8/3}$ -LQG surfaces

In this subsection we let $D \subset \mathbb{C}$ be a connected open set and we let h be a random distribution on D with the following property. For each bounded deterministic open set $U \subset D$ at positive Euclidean distance from ∂D ,

the law of $h|_U$ is absolutely continuous with respect to the corresponding restriction of an embedding into \mathbb{C} of a quantum sphere (with possibly random area). For example, h could be an embedding of a quantum disk, a thick ($\alpha \leq Q$) quantum wedge, a single bead of a thin ($\alpha \in (Q, Q + \sqrt{2/3})$) quantum wedge, or a quantum cone (in this last case we take D to be the complement of the two marked points for the cone). We will show how to obtain a $\sqrt{8/3}$ -LQG metric \mathfrak{d}_h on D from h .

The discussion at the end of Section 2.2.1 together with local absolute continuity allows us to define a metric $\mathfrak{d}_{h|_U}$ for any bounded open connected set $U \subset D$ at positive Euclidean distance from ∂D . If we let $\{U_n\}_{n \in \mathbb{N}}$ be an increasing sequence of such open sets with $\bigcup_{n=1}^{\infty} U_n = D$, then the metrics $\mathfrak{d}_{h|_{U_n}}$ (extended to be identically equal to ∞ for points outside of U_n) are decreasing as $n \rightarrow \infty$, so the limit

$$\mathfrak{d}_h(z, w) := \lim_{n \rightarrow \infty} \mathfrak{d}_{h|_{U_n}}(z, w)$$

exists for each $z, w \in D$ and defines a metric on D . It is easy to see that this metric does not depend on the choice of $\{U_n\}_{n \in \mathbb{N}}$.

We now record some elementary properties of the metric \mathfrak{d}_h . The first property is immediate from the above definition.

Lemma 2.2. *Suppose we are in the setting described just above, so that \mathfrak{d}_h is a well-defined metric on D . For any deterministic open connected set $U \subset \mathbb{C}$, the internal metric (Section 2.1) of \mathfrak{d}_h on U is a.s. equal to $\mathfrak{d}_{h|_U}$.*

Remark 2.3. It follows from Lemma 2.2 that if $A \subset \partial U$ and $\mathfrak{d}_{h|_U}$ extends by continuity to a metric on $U \cup A$ which induces the same topology on $U \cup A$ as the Euclidean metric (e.g., using the criterion of Lemma 2.5 just below) then for $x, y \in U \cup A$, we have $\mathfrak{d}_{h|_U}(x, y) \geq \mathfrak{d}_h(x, y)$. Indeed, the statement of the lemma immediately implies that this is the case whenever $x, y \in U$. In general, we take limits and use that both $\mathfrak{d}_{h|_{U \cup A}}$ and $\mathfrak{d}_h|_{U \cup A}$ (and the Euclidean metric) induce the same topology on U . We do *not* prove in this paper that $\mathfrak{d}_{h|_U}$ is the same as the internal metric of \mathfrak{d}_h on $U \cup A$.

Next we note that the LQG metric is coordinate invariant.

Lemma 2.4. *Suppose we are in the setting above. Let \tilde{D} be another domain and let $\phi : \tilde{D} \rightarrow D$ be a conformal map. If we let $\tilde{h} := h \circ \phi + Q \log |\phi'|$ (with Q as in (1.2)) then a.s. $\mathfrak{d}_h(\phi(z), \phi(w)) = \mathfrak{d}_{\tilde{h}}(z, w)$ for each $z, w \in \tilde{D}$.*

Proof. This follows since all of the quantities involved in the definition of the QLE(8/3, 0) processes $\tilde{\Gamma}^{z, w}$ used to define the metric are preserved under coordinate changes as in the statement of the lemma. \square

Finally, we give conditions under which \mathfrak{d}_h extends continuously to a subset of ∂D .

Lemma 2.5. *Suppose we are in the setting above and that D is simply connected, with simple boundary. Let $A \subset \partial D$ be a connected set and suppose that there is an open set $U \subset D$ such that $A \subset \partial U$, $\partial U \cap D$ lies at positive distance from A , and the law of $h|_U$ is absolutely continuous with respect to the law of the corresponding restriction of an embedding into D of a quantum disk (with fixed or random boundary length and area). Then \mathfrak{d}_h extends by continuity to a metric on $D \cup A$ which induces the Euclidean topology on $D \cup A$.*

In particular, suppose Q is as in (1.1), $x, y \in \partial D$, and h is either an embedding into (D, x, y) of an α -quantum wedge for $\alpha \leq Q$ or an embedding into (D, x, y) of a single bead of an α -quantum wedge with area $a > 0$ and left/right boundary lengths $\Gamma^-, \Gamma^+ > 0$ for $\alpha \in (Q, Q + \sqrt{2/3})$. Then a.s. \mathfrak{d}_h extends by continuity to $\partial D \setminus \{y\}$ (in the first case) or ∂D (in the second case).

Proof. In the case when h is in fact an embedding into D of a quantum disk, [MS16a, Corollary 1.5] implies that (D, \mathfrak{d}_h) is isometric to the Brownian disk. The Brownian disk metric extends continuously to the boundary, hence \mathfrak{d}_h does as well. In general, the first assertion of the lemma follows from local absolute continuity and the locality property of the $\sqrt{8/3}$ -LQG metric discussed above.

Since the laws of the particular choices of h discussed in the second statement are locally absolutely continuous with respect to the law of an embedding into D of a quantum disk away from x and y , it follows that the metric \mathfrak{d}_h in each of these cases extends by continuity to $D \setminus \{x, y\}$. Near x (and also near y , in the

case of a single bead of a thin wedge), the law of our given distribution h is locally absolutely continuous with respect to the law of an embedding of a β -quantum wedge into (D, x, y) for $\beta \leq Q$. Hence it suffices to show that if (D, h, x, y) is a β -quantum wedge for $\beta \leq Q$, then x a.s. lies at finite \mathfrak{d}_h -distance from the interior of D . By making a conformal change of coordinates and applying Lemma 2.4, we can assume without loss of generality that $(D, x, y) = (\mathbb{S}, \infty, -\infty)$ where $\mathbb{S} = \mathbb{R} \times (0, \pi)$ is the infinite horizontal strip.

In this case, [DMS14, Remark 4.5] implies that (after possibly applying a translation) the distribution h can be described as follows. If $\beta < Q$, let $X : \mathbb{R} \rightarrow \mathbb{R}$ be the process such that for $s \geq 0$, $X_s = B_{2s} - (Q - \beta)s$, where B is a standard linear Brownian motion with $B_0 = 0$; and for $s < 0$, $X_s = \widehat{B}_{-2s} + (Q - \beta)s$, where \widehat{B} is an independent standard linear Brownian motion with $\widehat{B}_0 = 0$ conditioned so that $\widehat{B}_{2t} + (Q - \beta)t > 0$ for each $t \geq 0$. If $\beta = Q$, instead let $X : \mathbb{R} \rightarrow \mathbb{R}$ be the process such that $\{X_s\}_{s \geq 0}$ is -1 times a 3-dimensional Bessel process started from 0 and $\{X_{-s}\}_{s \leq 0}$ is an independent standard linear Brownian motion started from 0. Then $h = h^0 + h^\dagger$, where h^0 is the function on \mathbb{S} such that $h^0(z) = X_s$ for $z \in \{s\} \times (0, \pi)$ for each $s \in \mathbb{R}$; and h^\dagger is a random distribution independent from h^0 whose law is that of the projection of a free boundary GFF on \mathbb{S} onto the space of functions on \mathbb{S} whose average over every segment $\{s\} \times (0, \pi)$ is 0.

By [MS16a, Proposition 6.7] (applied with $k = 0$), translation invariance of the law of h^\dagger , and the scaling property of the $\sqrt{8/3}$ -LQG metric [MS16a, Lemma 2.2], we infer that for each $k \in \mathbb{N}$,

$$\mathbb{E}[\mathfrak{d}_h((k-1) + i\pi/2, k + i\pi/2) \mid X] \leq C \exp\left(\frac{1}{\sqrt{6}} \sup_{s \in [k-1, k]} X_s\right).$$

for $C > 0$ a universal constant. Summing over all $k \in \mathbb{N}$ shows that a.s. $\mathfrak{d}_h(i\pi/2, \infty) < \infty$, as required. \square

3 The Brownian disk

For the proofs of our main results, we will require several facts about the Brownian disk, which was originally introduced in [BM15]. We collect these facts in this section.

3.1 Brownian disk definition

Fix $a, \ell > 0$. Here we will give the definition of the Brownian disk with area a and perimeter ℓ , following [BM15]. Let $X : [0, a] \rightarrow [0, \infty)$ be a standard Brownian motion started from ℓ and conditioned to hit 0 for the first time at time a (such a Brownian motion is defined rigorously in, e.g., [BM15, Section 2.1]). For $s, t \in [0, 1]$, set

$$d_X(s, t) := X_s + X_t - 2 \inf_{u \in [s \wedge t, s \vee t]} X_u. \quad (3.1)$$

As explained in [BM15, Section 2.1], d_X defines a pseudometric on $[0, a]$ and the quotient metric space $[0, 1]/\{d_X = 0\}$ is a forest of continuum random trees, indexed by the excursions of X away from its running infimum.

Conditioned on X , let Z^0 be the centered Gaussian process with

$$\text{Cov}(Z_s^0, Z_t^0) = \inf_{u \in [s \wedge t, s \vee t]} \left(X_u - \inf_{v \in [0, u]} X_v \right), \quad s, t \in [0, a].$$

One can readily check using the Kolmogorov continuity criterion that Z^0 a.s. admits a continuous modification which is α -Hölder continuous for each $\alpha < 1/4$. For this modification we have $Z_s^0 = Z_t^0$ whenever $d_X(s, t) = 0$, so Z^0 defines a function on the continuum random forest $[0, 1]/\{d_X = 0\}$.

Let \mathbf{b} be $\sqrt{3}$ times a Brownian bridge from 0 to 0 independent from (X, Z) with time duration ℓ . For $r \in [0, \ell]$, let $T_r := \inf\{t \geq 0 : X_t = \ell - r\}$ and for $t \in [0, a]$, let $T^{-1}(t) := \sup\{r \in [0, \ell] : T_r \leq t\}$. Set

$$Z_t := Z_t^0 + \mathbf{b}_{T^{-1}(t)}.$$

We view $[0, a]$ as a circle by identifying 0 with a and for $s, t \in [0, a]$ we define $\underline{Z}_{s,t}$ to be the minimal value of Z on the counterclockwise arc of $[0, a]$ from s to t . For $s, t \in [0, a]$, define

$$d_Z(s, t) = Z_s + Z_t - 2(\underline{Z}_{s,t} \vee \underline{Z}_{t,s}). \quad (3.2)$$

Then d_Z is not a pseudometric on $[0, a]$, so we define

$$d_{a,\ell}^0(s, t) = \inf \sum_{i=1}^k d_Z(s_i, t_i) \quad (3.3)$$

where the infimum is over all $k \in \mathbb{N}$ and all $2k+2$ -tuples $(t_0, s_1, t_1, \dots, s_k, t_k, s_{k+1}) \in [0, a]^{2k+2}$ with $t_0 = s$, $s_{k+1} = t$, and $d_X(t_{i-1}, s_i) = 0$ for each $i \in [1, k+1]_{\mathbb{Z}}$. In other words, $d_{a,\ell}^0$ is the largest pseudometric on $[0, a]$ which is at most d_Z and is zero whenever d_X is 0.

The *Brownian disk* with area a and perimeter ℓ is the quotient space $\text{BD}_{a,\ell} = [0, a]/\{d_{a,\ell}^0 = 0\}$ equipped with the quotient metric, which we call $d_{a,\ell}$. It is shown in [BM15] that $(\text{BD}_{a,\ell}, d_{a,\ell})$ is a.s. homeomorphic to the closed disk.

We write $\mathbf{p} : [0, a] \rightarrow \text{BD}_{a,\ell}$ for the quotient map. The pushforward $\mu_{a,\ell}$ of Lebesgue measure on $[0, a]$ under \mathbf{p} is a measure on $\text{BD}_{a,\ell}$ with total mass a , which we call the *area measure* of $\text{BD}_{a,\ell}$. The *boundary* of $\text{BD}_{a,\ell}$ is the set $\partial \text{BD}_{a,\ell} = \mathbf{p}(\{T_r : r \in [0, \ell]\})$. We note that $\text{BD}_{a,\ell}$ has a natural orientation, obtained by declaring that the path $t \mapsto \mathbf{p}(t)$ traces $\partial \text{BD}_{a,\ell}$ in the counterclockwise direction. The pushforward $\nu_{a,\ell}$ of Lebesgue measure on $[0, \ell]$ under the composition $r \mapsto \mathbf{p}(T_r)$ is called the *boundary measure* of $\text{BD}_{a,\ell}$.

By [MS16a, Corollary 1.5], the law of the metric measure space $(\text{BD}_{a,\ell}, d_{a,\ell}, \mu_{a,\ell}, \nu_{a,\ell})$ is the same as that of the $\sqrt{8/3}$ -LQG disk with area a and boundary length ℓ , equipped with its $\sqrt{8/3}$ -LQG area measure and boundary length measure.

Definition 3.1. Let $\ell > 0$ and let T_ℓ be a random variable with the law of the first time a standard linear Brownian motion hits $-\ell$. We write $(\text{BD}_{*,\ell}, d_{*,\ell}) := (\text{BD}_{T_\ell,\ell}, d_{T_\ell,\ell})$, so that $\text{BD}_{*,\ell}$ is a Brownian disk with random area. Note that in this case the corresponding function X defined above has the law of a standard linear Brownian motion started from ℓ and stopped at the first time it hits 0.

It is often convenient to work with a random-area Brownian disk rather than a fixed-area Brownian disk since the encoding function X has a simpler description in this case.

3.2 Area, length, and distance estimates for the Brownian disk

In this subsection we will prove some basic estimates relating distances, areas, and boundary lengths for the Brownian disk which are needed for the proofs of our main results. These estimates serve to quantify the intuition that for Brownian surfaces we have

$$\text{Area} \approx \text{Length}^{1/2} \approx \text{Distance}^{1/4}.$$

In other words, a subset of the Brownian disk with area δ typically has boundary length approximately $\delta^{1/2}$ and diameter approximately $\delta^{1/4}$. Our first estimate tells us that distances along the boundary are almost $1/2$ -Hölder continuous with respect to boundary length (in the same way that a standard Brownian motion on a compact time interval is almost $1/2$ -Hölder continuous).

Throughout this section, for $x, y \in \partial \text{BD}_{a,\ell}$ we write $[x, y]_{\partial \text{BD}_{a,\ell}}$ for the counterclockwise arc of $\partial \text{BD}_{a,\ell}$ from x to y (i.e., in the notation of Section 3.1, the arc traced by the boundary path $r \mapsto \mathbf{p}(T_r)$ between the times when it hits x and y).

Lemma 3.2. *Let $a > 0$ and $\ell > 0$ and let $(\text{BD}_{a,\ell}, d_{a,\ell})$ be a Brownian disk with area a and perimeter ℓ . For each $\zeta > 0$, there a.s. exists $C > 0$ such that for each $x, y \in \partial \text{BD}_{a,\ell}$, we have*

$$d_{a,\ell}(x, y) \leq C \nu_{a,\ell}([x, y]_{\partial \text{BD}_{a,\ell}})^{1/2} (|\log \nu_{a,\ell}([x, y]_{\partial \text{BD}_{a,\ell}})| + 1)^{\frac{7}{4} + \zeta} \quad (3.4)$$

where $[x, y]_{\partial \text{BD}_{a,\ell}}$ denotes the counterclockwise arc of $\partial \text{BD}_{a,\ell}$ from x to y . The same holds for a random-area disk with fixed boundary length as in Definition 3.1.

The idea of the proof of Lemma 3.2 is as follows. Recall that $\partial \text{BD}_{a,\ell}$ is the image under the quotient map $[0, a] \rightarrow \text{BD}_{a,\ell}$ of the set of times when the encoding function X attains a running infimum relative to time 0 (i.e. the image of $r \mapsto T_r$). The pair (X, Z^0) restricted to each such excursion evolves as a Brownian excursion together with the head of the Brownian snake driven by this excursion, so we can use tail bounds

for the Brownian snake [Ser97, Proposition 14] to bound the maximum of the restriction of Z^0 to each such excursion in terms of its time length. On the other hand, the excursions with unusually long time length are distributed according to a Poisson point process, so there cannot be too many such excursions in $[T_{r_1}, T_{r_2}]$ for any fixed $0 < r_1 < r_2 < \ell$. We use (3.3) to construct a path from the boundary points corresponding to T_{r_1} and T_{r_2} which skips all of the long excursions. See Figure 4 for an illustration. For the proof, we will use the elementary estimate

$$\mathbb{P}[X \geq x] \leq \frac{e^{-\lambda}(e\lambda)^x}{x^x}, \quad \forall x > \lambda \quad \text{for } X \sim \text{Poisson}(\lambda). \quad (3.5)$$




Figure 4: Illustration of the proof of Lemma 3.2. Shown is the Brownian disk $\text{BD}_{*,\ell}$ and an arc $\mathbb{P}(T([r_1, r_2])) \subset \partial \text{BD}_{*,\ell}$ (red) whose diameter we are trying to bound. To construct a path from $\mathbb{P}(T_{r_1})$ to $\mathbb{P}(T_{r_2})$, we consider a path (shown in blue) consisting of a concatenation of segments which avoid the large excursions of $t \mapsto \mathbb{P}(t)$ away from $\mathbb{P}(T([r_1, r_2]))$ (pink). The number of such excursions is bounded using (3.6) and the lengths of each of the segments between excursions is bounded using (3.2) and (3.12).

Proof of Lemma 3.2. First fix $\ell \in \mathbb{N}$ and let $\text{BD}_{*,\ell}$ be a random-area Brownian disk with boundary length ℓ . We use the notation introduced at the beginning of this subsection with $a = T_\ell$ as in Definition 3.1. For $r > 0$, let $\tilde{T}_r := \sup_{s < r} T_s$. Note that $d_X(\tilde{T}_r, T_r) = 0$, with d_X as in (3.1). Let \mathcal{R} be the set of $r \in [0, \ell]$ for which $\tilde{T}_r \neq T_r$, so that the intervals $[\tilde{T}_r, T_r]$ for $r \in \mathcal{R}$ are precisely the excursion intervals for X away from its running infimum.

The time lengths of the excursions of X away from its running infimum, parameterized by minus the running infimum of X , have the law of a Poisson point process on \mathbb{R} with Itô measure $\pi^{-1/2}t^{-3/2}dt$ (see, e.g. [RY99, Theorem 2.4 and Proposition 2.8, Section XII]). Hence for $0 \leq r_1 \leq r_2 \leq \ell$ and $A > 0$, the law of $\#\{r \in [r_1, r_2] : T_r - \tilde{T}_r > A\}$ is Poisson with mean $2\pi^{-1/2}(r_2 - r_1)A^{-1/2}$. By (3.5) and the union bound, for each $k \in \mathbb{N}$ it holds except on an event of probability decaying faster than any exponential function of k that

$$\#\{r \in [(j-1)2^{-k}, j2^{-k}] : T_r - \tilde{T}_r > 2^{-2k}\} \leq k, \quad \forall j \in [1, 2^k]_{\mathbb{Z}}. \quad (3.6)$$

For $n \in \mathbb{N}$, let \mathcal{R}_n be the set of $r \in [0, \ell]$ for which $T_r - \tilde{T}_r \in [(n+1)^{-1}\ell^2, n^{-1}\ell^2]$. Then $\#\mathcal{R}_n$ is Poisson with mean $2\pi^{-1/2}((n+1)^{1/2} - n^{1/2}) \asymp n^{-1/2}$. Hence, by (3.5), except on an event of probability decaying

faster than any power of n , we have

$$\#\mathcal{R}_m \leq \log(m+1), \quad \forall m \geq n. \quad (3.7)$$

Let \mathcal{F} be the σ -algebra generated by the set \mathcal{R} and the pairs (\tilde{T}_r, T_r) for $r \in \mathcal{R}$. If we condition on \mathcal{F} then the conditional law of the processes $\{(X - X_{\tilde{T}_r}, Z^0)|_{[\tilde{T}_r, T_r]} : r \in \mathcal{R}\}$ is described as follows. These processes for different choices of r are conditionally independent given \mathcal{F} ; the conditional law of each $(X - X_{\tilde{T}_r})|_{[\tilde{T}_r, T_r]}$ is that of a Brownian excursion with time length $T_r - \tilde{T}_r$; and each $Z^0|_{[\tilde{T}_r, T_r]}$ is the head of the Brownian snake driven by $(X - X_{\tilde{T}_r})|_{[\tilde{T}_r, T_r]}$. By the large deviation estimate for the head of a Brownian snake driven by a standard Brownian excursion [Ser97, Proposition 14] and two applications of Brownian scaling, we find that for each $r \in \mathcal{R}$ and each $A > 1$,

$$\mathbb{P} \left[\sup_{t \in [\tilde{T}_r, T_r]} |Z_s^0 - Z_t^0| > A(T_r - \tilde{T}_r)^{1/4} \mid \mathcal{F} \right] \leq \exp \left(-\frac{3}{2}(1 + o_A(1))A^{4/3} \right) \quad (3.8)$$

at a universal rate as $A \rightarrow \infty$.

If the event in (3.7) occurs for some $n \in \mathbb{N}$ (which we emphasize is \mathcal{F} -measurable), then (3.8) implies that with

$$E_n := \left\{ \exists r \in \bigcup_{m=n}^{\infty} \mathcal{R}_m \text{ with } \sup_{t \in [\tilde{T}_r, T_r]} |Z_t^0| > \ell^{1/2} n^{-1/4} \log(n)^{\frac{3}{4} + \zeta} \right\} \quad (3.9)$$

we have

$$\mathbb{P}[E_n \mid \mathcal{F}] \leq \sum_{m=n}^{\infty} \log(m+1) \exp \left(-\frac{3}{2}(1 + o_n(1)) \log(n)^{1 + \frac{4}{3}\zeta} (m/n)^{1/3} \right) = o_n(n^{-p}), \quad \forall p > 0. \quad (3.10)$$

Recall that $Z = Z^0 + \mathbf{b}_{T^{-1}(\cdot)}$, where \mathbf{b} is equal to $\sqrt{3}$ times a reparameterized Brownian bridge from 0 to 0 in time ℓ . By a straightforward Gaussian estimate, it holds except on an event of probability decaying faster than any power of n that whenever $0 \leq r_1 \leq r_2 \leq \ell$ with $r_2 - r_1 \leq \ell n^{-1/2}$ we have

$$\sup_{\rho_1, \rho_2 \in [r_1, r_2]} |\mathbf{b}_{\rho_1} - \mathbf{b}_{\rho_2}| \leq \ell^{1/2} n^{-1/4} \log(n)^{3/4}. \quad (3.11)$$

If the event E_n in (3.9) does not occur, then for any such $r_1, r_2 \in [0, \ell]$ and any $t \in [T_{r_1}, T_{r_2}]$ for which $t \notin \bigcup_{m=1}^{n-1} \bigcup_{r \in \mathcal{R}_m} [\tilde{T}_r, T_r]$, we have either $Z_t^0 = 0$ or $t \in [\tilde{T}_r, T_r]$ for some $r \in \bigcup_{m=n}^{\infty} \mathcal{R}_m$. In either case $|Z_t^0| \leq \ell^{1/2} n^{-1/4} \log(n)^{3/4 + \zeta}$. Hence (3.10), (3.11), and the triangle inequality together imply that except on an event of probability decaying faster than any power of n ,

$$\sup \left\{ |Z_s - Z_t| : s, t \in [T_{r_1}, T_{r_2}] \setminus \bigcup_{m=1}^{n-1} \bigcup_{r \in \mathcal{R}_m} [\tilde{T}_r, T_r] \right\} \leq 4\ell^{1/2} n^{-1/4} \log(n)^{\frac{3}{4} + \zeta} \quad (3.12)$$

simultaneously for each $0 \leq r_1 < r_2 \leq \ell$ with $r_2 - r_1 \leq \ell n^{-1/2}$.

Suppose now that $k \in \mathbb{N}$, (3.6) occurs and (3.12) occurs with $n = 2^{2k}$. Let $\mathbf{p} : [0, T_\ell] \rightarrow \text{BD}_{*, \ell}$ be the quotient map. Note that $\mathbf{p}(T([(j-1)2^{-k}\ell, j2^{-k}\ell]))$ corresponds to the counterclockwise segment of the boundary of the disk of length $2^{-k}\ell$ which connects $\mathbf{p}(T_{(j-1)2^{-k}\ell})$ to $\mathbf{p}(T_{j2^{-k}\ell})$. We will use (3.3) to bound the diameter of $\mathbf{p}(T([(j-1)2^{-k}\ell, j2^{-k}\ell]))$. Fix $j \in [1, 2^k]_{\mathbb{Z}}$ and $r_1, r_2 \in [(j-1)2^{-k}\ell, j2^{-k}\ell]$ with $r_1 < r_2$. Let $t_0 = s_1 = T_{r_1}$, $t_k = s_{k+1} = T_{r_2}$, and for $i \in [2, k]_{\mathbb{Z}}$ let $[t_{i-1}, s_i]$ be the i th interval $[\tilde{T}_r, T_r] \subset [T_{r_1}, T_{r_2}]$ with $T_r - \tilde{T}_r \geq \ell 2^{-2k}$, counted from left to right; or $t_{i-1} = s_i = T_{r_2}$ if there are fewer than i such intervals. Then we have $d_X(t_{i-1}, s_i) = 0$ for each $i \in [1, k+1]_{\mathbb{Z}}$.

By (3.6) each $r \in [r_1, r_2] \cap \bigcup_{m=1}^{2^{2k}} \mathcal{R}_m$ satisfies $[\tilde{T}_r, T_r] = [t_{i-1}, s_i]$ for some $i \in [2, k]_{\mathbb{Z}}$. Hence each of the intervals $[s_i, t_i]$ for $i \in [1, k]_{\mathbb{Z}}$ is disjoint from $\bigcup_{m=1}^{2^{2k}} \bigcup_{r \in \mathcal{R}_m} [\tilde{T}_r, T_r]$. By (3.3) and (3.12),

$$d_{*, \ell}^0(T_{r_1}, T_{r_2}) \leq 2 \sum_{i=1}^k \sup_{s, t \in [s_i, t_i]} |Z_s - Z_t| \leq \ell^{1/2} 2^{-k/2} k^{\frac{7}{4} + \zeta},$$

with universal implicit constant, where in the first inequality we recall the definition (3.2) of d_Z . Since our choice of $r_1, r_2 \in [(j-1)2^{-k}, j2^{-k}]$ was arbitrary,

$$\text{diam}(\mathbb{p}(T([(j-1)2^{-k}\ell, j2^{-k}\ell])); d_{*,\ell}) \preceq \ell^{1/2} 2^{-k/2} k^{\frac{7}{4}+\zeta} \quad (3.13)$$

where here we recall the construction of $d_{*,\ell}$ from $d_{*,\ell}^0$.

By the Borel-Cantelli lemma, there a.s. exists $k_0 \in \mathbb{N}$ such that for $k \geq k_0$, (3.6) occurs and (3.12) occurs with $n = 2^{2k}$. Suppose given $k \geq k_0$ and $x, y \in \partial \text{BD}_{*,\ell}$ with $2^{-k-1} \leq \nu_{*,\ell}([x, y]_{\partial \text{BD}_{*,\ell}}) \leq 2^{-k}$. By (3.13) and the triangle inequality,

$$d_{*,\ell}(x, y) \leq C \nu_{*,\ell}([x, y]_{\partial \text{BD}_{*,\ell}})^{1/2} (|\log \nu_{*,\ell}([x, y]_{\partial \text{BD}_{*,\ell}})| + 1)^{\frac{7}{4}+\zeta}$$

for an appropriate $C > 0$ as in the statement of the lemma. By increasing the constant to deal with the case when $\nu_{*,\ell}([x, y]_{\partial \text{BD}_{*,\ell}}) > 2^{-k_0}$, we get the desired continuity estimate in the case of a random-area disk.

The fixed area case follows since for any $a > 0$, the law of $(\text{BD}_{a,\ell}, d_{a,\ell})$ is absolutely continuous with respect to the law of $(\text{BD}_{*,2\ell}, d_{*,2\ell})$ conditioned on the positive probability event that $T_\ell > 2a$ (c.f. [BM15, Section 2.1]). \square

Next we need an estimate for the areas of metric balls, which is a straightforward consequence of the Hölder continuity of Z and the upper bound for areas of metric balls in the Brownian map [Le 10, Corollary 6.2].

Lemma 3.3. *Let $a, \ell > 0$ and let $(\text{BD}_{a,\ell}, d_{a,\ell})$ be a Brownian disk with area a and perimeter ℓ . For each $u \in (0, 1)$, there a.s. exists $C > 1$ such that the following is true. For each $\delta \in (0, 1)$ and each $z \in \text{BD}_{a,\ell}$, we have*

$$\mu_{a,\ell}(B_\delta(z; d_{a,\ell})) \geq C^{-1} \delta^{4+u} \quad (3.14)$$

and for each $z \in \text{BD}_{a,\ell}$ with $B_\delta(z; d_{a,\ell}) \cap \partial \text{BD}_{a,\ell} = \emptyset$, we have

$$\mu_{a,\ell}(B_\delta(z; d_{a,\ell})) \leq C \delta^{4-u}. \quad (3.15)$$

Proof. It is easy to see from the Kolmogorov continuity criterion that the process Z used in the definition of the Brownian disk is a.s. Hölder continuous with exponent $(4+u)^{-1}$.

It follows from (3.2) that there a.s. exists a random $C > 0$ such that for each $s, t \in [0, a]$, we have

$$d_{a,\ell}(\mathbb{p}(s), \mathbb{p}(t)) \leq C |t - s|^{(4+u)^{-1}}. \quad (3.16)$$

Since the quotient map \mathbb{p} pushes forward Lebesgue measure on $[0, a]$ to $\mu_{a,\ell}$, we infer that for each $t \in [0, \ell]$ and each sufficiently small $\delta \in (0, 1)$ (how small depends on C, u , and a),

$$\mu_{a,\ell}(B_\delta(\mathbb{p}(t); d_{a,\ell})) \geq C^{-1} \delta^{4+u}.$$

Upon shrinking C , this implies (3.14).

We now prove (3.15). By local absolute continuity of the process (Z, X, \mathbf{b}) for different choices of a and ℓ , it suffices to prove that for Lebesgue a.e. pair $(a, \ell) \in (0, \infty)^2$, there a.s. exists $C > 0$ so that (3.15) holds. Suppose to the contrary that this is not the case. Then there is a positive Lebesgue measure set $\mathcal{A} \subset (0, \infty)^2$ such that for each $(a, \ell) \in \mathcal{A}$, it holds with positive probability that

$$\sup_{\delta > 0} \sup_{z \in \text{BD}_{a,\ell}} \frac{\mu_{a,\ell}(B_\delta(z; d_{a,\ell}))}{\delta^{4-u}} = \infty. \quad (3.17)$$

Choose $A > 0$ such that the projection of \mathcal{A} onto its first coordinate intersects $[0, A]$ in a set of positive Lebesgue measure and let (BM_A, d_A) be a Brownian map with area A and let μ_A be its area measure. By [Le 10, Corollary 6.2], a.s.

$$\sup_{\delta > 0} \sup_{z \in \text{BM}_A} \frac{\mu_A(B_\delta(z; d_A))}{\delta^{4-u}} < \infty. \quad (3.18)$$

Now fix $r > 0$ and let $z_0, z_1 \in \text{BM}_A$ be sampled uniformly from μ_A . By [MS15a, Proposition 4.4], the complementary connected component D of $B_r(z_0; d_A)$ containing z_1 , equipped with the internal metric $d_{A,D}$ induced by d_A , has the law of a Brownian disk if we condition on its area and boundary length. With positive probability, the area and boundary length of D belong to \mathcal{A} . Since each $d_{A,D}$ -metric ball is contained in a d_A -metric ball with the same radius, we see that (3.17) contradicts (3.18). \square

We will next prove a lower bound for the amount of area near a boundary arc of given length.

Lemma 3.4. *Let $a, \ell > 0$ and let $(\text{BD}_{a,\ell}, d_{a,\ell})$ be a Brownian disk with area a and perimeter ℓ . For each $u > 0$, there a.s. exists $c > 0$ such that for each $\delta \in (0, 1)$ and each $x, y \in \partial \text{BD}_{a,\ell}$, we have*

$$\mu_{a,\ell}(B_\delta([x, y]_{\partial \text{BD}_{a,\ell}}; d_{a,\ell})) \geq c\delta^{2+u}\nu_{a,\ell}([x, y]_{\partial \text{BD}_{a,\ell}}).$$

The idea of the proof of Lemma 3.4 is to prove a lower bound for the number of “large” excursions of X away from its running infimum during the time interval $[T_{r_1}, T_{r_2}]$ where $0 \leq r_1 < r_2 < \ell$ are chosen so that the times T_{r_1} and T_{r_2} are mapped to x and y under the quotient map $[0, a] \rightarrow \text{BD}_{a,\ell}$. This lower bound complements the upper bound for the number of such excursions used in the proof of Lemma 3.2. For the proof of Lemma 3.4 we will need the following elementary estimate for Brownian motion.

Lemma 3.5. *Let B be a standard linear Brownian motion started from 0 and for $r > 0$, let $T_r := \inf\{t > 0 : B_t = -r\}$. For $\delta > 0$, let N_δ be the number of intervals of the form $[(k-1)\delta, k\delta]$ for $k \in \mathbb{N}$ which intersect $\{T_r : r \in [0, 1]\}$. There is a universal constant $c_0 > 0$ such that for each $\delta \in (0, 1)$ and each $\zeta \in (0, (2\pi)^{-1/2})$ we have*

$$\mathbb{P}[N_\delta < \zeta\delta^{-1/2}] \leq \exp\left(-\frac{c_0}{\zeta\delta^{1/2}}\right).$$

Proof. Let $\rho_0 = 0$ and for $j \in \mathbb{N}$ inductively let ρ_j be the smallest $r > \rho_{j-1}$ for which $T_r - T_{\rho_{j-1}} \geq \delta$. Let J be the largest $j \in \mathbb{N}$ for which $\rho_j \leq 1$. Each of the times T_{ρ_j} for $j \in [1, J]_{\mathbb{Z}}$ lies in a distinct interval of the form $[(k-1)\delta, k\delta]$ for $k \in \mathbb{N}$. Hence $N_\delta \geq J$.

By the strong Markov property for standard Brownian motion, the random variables $\rho_j - \rho_{j-1}$ for $j \in \mathbb{N}$ are i.i.d. Observe that $\rho_1 - \rho_0 = -\inf_{t \in [0, \delta]} B_t$ because if $r < -\inf_{t \in [0, \delta]} B_t$ then $T_r < \delta$ and if $r > -\inf_{t \in [0, \delta]} B_t$ then $T_r > \delta$. Consequently, it follows that each of these random variables has the law of the absolute value of a Gaussian random variable with mean 0 and variance δ (which has mean $(2/\pi)^{1/2}\delta^{1/2}$). By Hoeffding’s inequality for sums of independent random variables with sub-Gaussian tails (see, e.g., [Ver12, Proposition 5.10]), for $m \in \mathbb{N}$ and $R > 0$,

$$\mathbb{P}[\rho_m > R + (2/\pi)^{1/2}\delta^{1/2}m] \leq \exp\left(-\frac{c_1 R^2}{\delta m}\right)$$

with $c_1 > 0$ a universal constant. Therefore, for $\zeta \in (0, (2\pi)^{-1/2})$ we have

$$\mathbb{P}[N_\delta < \zeta\delta^{-1/2}] \leq \mathbb{P}[J < \zeta\delta^{-1/2}] = \mathbb{P}[\rho_{\lfloor \zeta\delta^{-1/2} \rfloor} > 1] \leq \exp\left(-\frac{c_0}{\zeta\delta^{1/2}}\right)$$

with $c_0 > 0$ as in the statement of the lemma. \square

Proof of Lemma 3.4. In light of Lemma 3.3 (applied with $u/2$ in place of u), we can restrict attention to arcs $[x, y]_{\partial \text{BD}_{a,\ell}}$ satisfying

$$\nu_{a,\ell}([x, y]_{\partial \text{BD}_{a,\ell}}) \geq \delta^{2-u/2}.$$

We start by working with a random-area Brownian disk $(\text{BD}_{*,\ell}, d_{*,\ell})$ as in Definition 3.1. Fix a small parameter $v \in (0, 1)$ to be chosen later, depending only on u . For $n, m \in \mathbb{N}$ with $n \geq m$ and $j \in [1, 2^m]_{\mathbb{Z}}$, let $N_{m,j}^n$ be the number of intervals of the form $[(k-1)2^{-2n}, k2^{-2n}]$ for $k \in \mathbb{N}$ which intersect $\{T_r : r \in [(j-1)2^{-m}\ell, j2^{-m}\ell]\}$. Let E_m be the event that $N_{m,j}^n \geq \lfloor \ell 2^{(1-v)n-m} \rfloor$ for each $n \geq m$ and all $j \in [1, 2^m]_{\mathbb{Z}}$. By Lemma 3.5 (applied with $\delta = 2^{2m-2n}\ell^{-2}$ and $\zeta = 2^{-vn}$) and scale and translation invariance,

$$\mathbb{P}[N_{m,j}^n < \lfloor \ell 2^{(1-v)n-m} \rfloor] \leq \exp\left(-c_0 \ell 2^{(1+v)n-m}\right).$$

By the union bound,

$$\mathbb{P}[E_m^c] \leq 2^m \sum_{n=m}^{\infty} \exp\left(-c_0 \ell 2^{(1+v)n-m}\right) \leq 2^m \exp(-c_0 \ell 2^{vm})$$

with implicit constant depending only on ℓ . By the Borel-Cantelli Lemma, a.s. there exists $m_0 \in \mathbb{N}$ such that E_m occurs for each $m \geq m_0$.

Let $C > 0$ be a random constant chosen so that (3.16) holds for $\text{BD}_{*,\ell}$, with v in place of u . Suppose we are given $\delta > 0$ with $\delta^2 \in (0, 2^{-(1+2v)m_0}]$ and $x, y \in \partial \text{BD}_{a,\ell}$ with

$$\delta^{\frac{2}{1+2v}} \leq \nu_{a,\ell}([x, y]_{\partial \text{BD}_{a,\ell}}) \leq 2^{-m_0-1}. \quad (3.19)$$

After possibly increasing m_0 , we can choose $m \geq m_0$ with $2^{-m+1} \leq \nu_{*,\ell}([x, y]_{\partial \text{BD}_{*,\ell}}) \leq 2^{-m+2}$ and $n \geq m$ with $2^{-2(1+v)n} \leq C^{-4}\delta^4 \leq 2^{-2(1+v)n+1}$.

For some $j \in [1, 2^m]_{\mathbb{Z}}$, the boundary arc $[x, y]_{\partial \text{BD}_{*,\ell}}$ contains the image under the quotient map \mathfrak{p} of the set $\{T_r : r \in [(j-1)2^{-m}\ell, j2^{-m}\ell]\}$. By definition of E_m , there are at least $\lfloor \ell 2^{(1-v)n-m} \rfloor$ intervals of the form $[(k-1)2^{-2n}, k2^{-2n}]$ for $k \in [1, \ell 2^{2n}]_{\mathbb{Z}}$ whose images under \mathfrak{p} intersect $[x, y]_{\partial \text{BD}_{*,\ell}}$. By (3.16) and our choice of n , if v is chosen sufficiently small relative to u (depending on the constant C from (3.16)) then the image of each of these intervals under \mathfrak{p} lies in $B_\delta([x, y]_{\partial \text{BD}_{*,\ell}}; d_{*,\ell})$. Therefore, for an appropriate random choice of $c' > c > 0$,

$$\mu_{*,\ell}(B_\delta([x, y]_{\partial \text{BD}_{*,\ell}}; d_{*,\ell})) \geq c' 2^{-(1+v)n-m} \geq c\delta^2 \nu_{*,\ell}([x, y]_{\partial \text{BD}_{*,\ell}}).$$

This holds simultaneously for each boundary arc $[x, y]_{\partial \text{BD}_{*,\ell}}$ satisfying (3.19). By choosing v sufficiently small relative to u (recall (3.16)) and shrinking c in a manner depending on m_0 , we obtain the statement of the lemma for a random-area Brownian disk. The statement for a fixed area Brownian disk follows by local absolute continuity. \square

4 Metric gluing

In this section we will complete the proofs of the theorems stated in Section 1.3. In Section 4.1, we will show that $\sqrt{8/3}$ -LQG geodesics between quantum typical points a.s. do not hit the boundary of the domain, which is one of the main inputs in the proofs of our main results. Next, in Section 4.2, we will prove Theorem 1.1, noting that the proof of Theorem 1.3 is essentially identical. Finally, in Section 4.3, we will deduce Theorems 1.6 and 1.7.

Throughout this section, we use the notation

$$\mathbb{V}_\rho := \{z \in B_\rho(0) : \text{Im } z > \rho^{-1}\} \quad \text{and} \quad \mathbb{V}'_\rho := \{z \in B_\rho(0) : |z| > \rho^{-1}\}, \quad \forall \rho > 1. \quad (4.1)$$

4.1 LQG geodesics cannot hit the boundary

To prove our main results, we want to apply the estimates for the Brownian disk obtained in Section 3.2. In order to apply these estimates, we need to ensure that we can restrict attention to finitely many quantum surfaces that locally look like quantum disks (equivalently, Brownian disks).

If we are in the setting of Theorem 1.1 with either $\mathfrak{w}^- < 4/3$ or $\mathfrak{w}^+ < 4/3$, the $\text{SLE}_{8/3}(\mathfrak{w}^- - 2; \mathfrak{w}^+ - 2)$ curve η will intersect \mathbb{R} in a fractal set (see [MW14] for a computation of the dimension of this set), so there will be infinitely many elements of $\mathcal{U}^- \cup \mathcal{U}^+$ (i.e., connected components of $\mathbb{H} \setminus \eta$) contained in small neighborhoods of certain points of \mathbb{R} . However, since η does not intersect itself and is a.s. continuous and transient, there are only finitely many connected components of $\mathbb{H} \setminus \eta$ which intersect each of the sets \mathbb{V}_ρ of (4.1). Hence one way to avoid dealing with infinitely many elements of $\mathcal{U}^- \cup \mathcal{U}^+$ is to work in a bounded set at positive distance from \mathbb{R} . The following lemma will allow us to do so. For the statement, we recall the parameter Q from (1.2).

Lemma 4.1. *Suppose that h is either a free boundary GFF on \mathbb{H} plus $-\alpha \log |\cdot|$ for $\alpha \leq Q$, an embedding into $(\mathbb{H}, 0, \infty)$ of an α -quantum wedge for $\alpha \leq Q$, or an embedding into $(\mathbb{H}, 0, \infty)$ of a single bead of an α -quantum wedge for $\alpha \in (Q, Q + \sqrt{2/3})$ with area $\mathfrak{a} > 0$ and left/right boundary lengths $\mathfrak{l}^-, \mathfrak{l}^+ > 0$. Let $R > 1$ and let z_1, z_2 be sampled uniformly from $\mu_h|_{\mathbb{V}_R}$, normalized to be a probability measure, where \mathbb{V}_R is as in (4.1). Almost surely, there is a unique \mathfrak{d}_h -geodesic γ_{z_1, z_2} from z_1 to z_2 , and a.s. γ_{z_1, z_2} does not intersect $\mathbb{R} \cup \{\infty\}$.*

We will eventually deduce Lemma 4.1 from [BM15, Lemma 18] and local absolute continuity of h with respect to an embedding of a quantum disk (which is isometric to a Brownian disk). However, due to the presence of the log singularity at 0 this absolute continuity only holds away from 0 and ∞ so we first need to rule out the possibility that \mathfrak{d}_h -geodesics hit 0 or ∞ with positive probability. By an absolute continuity argument, it suffices to prove this in the case when h is a free boundary GFF with a log singularity at 0. This is the purpose of the next two lemmas.

The proof of the following lemma illustrates a general technique which can be used to show that various events associated with the $\sqrt{8/3}$ -LQG metric induced by some variant of the GFF occur with positive probability.

Lemma 4.2. *Let h be a free-boundary GFF on a simply connected domain $D \subset \mathbb{C}$. Let $A_1, A_2 \subset \overline{D}$ be deterministic disjoint compact sets and let $c > 0$. Then with positive probability, the $\sqrt{8/3}$ -LQG metric \mathfrak{d}_h satisfies*

$$\text{diam}_{\mathfrak{d}_h}(A_1) \leq c \mathfrak{d}_h(A_1, A_2). \quad (4.2)$$

Proof. Let $U_1, U_2 \subset \overline{D}$ be bounded connected relatively open sets such that $A_1 \subset U_1$, $A_2 \subset U_2$, and $\overline{U_1} \cap \overline{U_2} = \emptyset$. We note that $h|_{U_1}$ has the law of a GFF on U_1 plus a random harmonic function. Hence the $\sqrt{8/3}$ -LQG metric $\mathfrak{d}_{h|_{U_1}}$ is well-defined, finite on compact subsets of U_1 , and determines the same topology on U_1 as the Euclidean metric. We furthermore have $\mathfrak{d}_{h|_{U_1}} \geq \mathfrak{d}_h$ on U_1 . Since \mathfrak{d}_h determines the same topology as the Euclidean metric, there exists $C > 0$ such that

$$\mathbb{P}\left[\text{diam}\left(A_1; \mathfrak{d}_{h|_{U_1}}\right) \leq C \mathfrak{d}_h(A_2, \partial U_2)\right] \geq \frac{1}{2}. \quad (4.3)$$

Let $R > 0$ be a constant to be chosen later and let ϕ be a smooth function on \overline{D} which is identically equal to $-R$ on U_1 , identically equal to R on U_2 , and identically equal to 0 outside a compact subset of \overline{D} . Let $\hat{h} := h + \phi$. By the scaling property of the $\sqrt{8/3}$ -LQG metric [MS16a, Lemma 2.2],

$$\text{diam}\left(A_1; \mathfrak{d}_{\hat{h}|_{U_1}}\right) = e^{-\sqrt{8/3}R/4} \text{diam}\left(A_1; \mathfrak{d}_{h|_{U_1}}\right) \quad \text{and} \quad \mathfrak{d}_{\hat{h}}(A_2, \partial U_2) = e^{\sqrt{8/3}R/4} \mathfrak{d}_h(A_2, \partial U_2).$$

Hence if we choose $R > 0$ such that $Ce^{-\sqrt{8/3}R/4} \leq c$, then (4.3) implies that with probability at least $1/2$,

$$\text{diam}\left(A_1; \mathfrak{d}_{\hat{h}|_{U_1}}\right) \leq c \mathfrak{d}_{\hat{h}}(A_2, \partial U_2),$$

in which case

$$\text{diam}(A_1; \mathfrak{d}_{\hat{h}}) \leq c \mathfrak{d}_{\hat{h}}(A_1, A_2). \quad (4.4)$$

On the other hand, since $\hat{h} - h$ is a smooth compactly supported function, the law of \hat{h} is absolutely continuous with respect to the law of h , so with positive probability (4.4) holds with h in place of \hat{h} . By scaling, this implies that (4.4) holds. \square

Now we can rule out the possibility that geodesics for the $\sqrt{8/3}$ -LQG metric induced by a GFF with a log singularity at 0 hit 0.

Lemma 4.3. *Let \tilde{h} be a free boundary GFF on \mathbb{H} and let $\alpha \in \mathbb{R}$. Let $h := \tilde{h} - \alpha \log|\cdot|$ and let \mathfrak{d}_h be the $\sqrt{8/3}$ -LQG metric induced by h . Almost surely, no \mathfrak{d}_h -geodesic between points in \mathbb{H} hits 0.*

Proof. For each $j \in \mathbb{N}$, let $r_j = e^{-j}$ and let \mathcal{F}_j be the σ -algebra generated by $h|_{\mathbb{H} \setminus B_{r_j}(0)}$. Let \mathfrak{h}_j be the conditional mean of $h|_{B_{r_j}(0) \cap \mathbb{H}}$ given $h|_{\mathbb{H} \setminus B_{r_j}(0)}$. Let E_j be the event that

$$\text{diam}(\partial B_{r_{j-1}}(0) \cap \mathbb{H}; \mathfrak{d}_h) < \mathfrak{d}_h(\partial B_{r_{j-1}}(0) \cap \mathbb{H}, (\partial B_{r_j}(0) \cup \partial B_{r_{j-2}}(0)) \cap \mathbb{H}).$$

Then if E_j occurs for some $j \in \mathbb{N}$, no geodesic between points in $\mathbb{H} \setminus B_{r_j}(0)$ can hit 0. Hence we just need to show that for each $j_0 \in \mathbb{N}$,

$$\mathbb{P}\left[\bigcup_{j=j_0}^{\infty} E_j\right] = 1.$$

The event E_j is the same as the event that the following is true: if we grow the \mathfrak{d}_h -metric balls centered at any point of $\partial B_{r_{j-1}}(0)$, then we cover $\partial B_{r_{j-1}}(0)$ before reaching $\partial B_{r_j}(0)$ or $\partial B_{r_{j-2}}(0)$. Since metric balls are locally determined by h (this is immediate from the construction of \mathfrak{d}_h using QLE(8/3, 0) in [MS15b, MS16a, MS16b]), it follows that E_j is determined by the restriction of h to $B_{r_{j-2}}(0) \setminus B_{r_j}(0)$, and in particular $E_j \in \mathcal{F}_j$. By the scale invariance of \mathfrak{d}_h , the events E_j do not depend on the choice of additive constant for h . By the conformal invariance of the law of h , modulo additive constant, and Lemma 4.2, we can find $p > 0$ such that $\mathbb{P}[E_j] \geq p$ for each $j \in \mathbb{N}$.

Now fix $j_0 \in \mathbb{N}$ and $\epsilon > 0$. Inductively, suppose $k \in \mathbb{N}$ and we have defined a $j_{k-1} \geq j_0$ which is a stopping time for the filtration $\{\mathcal{F}_j\}_{j \in \mathbb{N}}$. The function $\mathfrak{h}_{j_{k-1}}$ is a.s. harmonic, hence smooth, on $\overline{B_{r_{j_{k-1}+1}}(0)} \cap \mathbb{H}$ so we can find $j_k \geq j_{k-1} + 2$ such that

$$\int_{B_{r_{j_k-2}}(0) \cap \mathbb{H}} |\nabla \mathfrak{h}_{j_{k-1}}(w)|^2 dw \leq \epsilon.$$

Since $\mathfrak{h}_{j_{k-1}}$ is $\mathcal{F}_{j_{k-1}}$ -measurable, it follows by induction that j_k is a stopping time for $\{\mathcal{F}_j\}_{j \in \mathbb{N}}$. By [MS16a, Lemma 5.4], if we choose $\epsilon > 0$ sufficiently small, depending on p , then it holds with conditional probability at least $1 - p/2$ given $\mathcal{F}_{j_{k-1}}$ that the Radon-Nikodym derivative of the conditional of $h|_{B_{r_{j_k-2}}(0) \cap \mathbb{H}}$ with respect to the unconditional law of a free-boundary GFF on \mathbb{H} plus $-\alpha \log |\cdot|$ restricted to $B_{r_{j_k-2}}(0) \cap \mathbb{H}$, viewed as distributions defined modulo additive constant, is between $1/2$ and 2 . Hence

$$\mathbb{P}[E_{j_k} | \mathcal{F}_{j_{k-1}}] \geq \frac{p}{4}.$$

By the Borel-Cantelli lemma, it follows that a.s. E_{j_k} occurs for some $k \in \mathbb{N}$. \square

Proof of Lemma 4.1. For each sufficiently small $r > 0$, the law of $h|_{B_r(0) \cap \mathbb{H}}$ for an appropriate choice of embedding h is absolutely continuous with respect to the law of the corresponding restriction of a free-boundary GFF plus a log singularity at 0 with appropriate choice of additive constant. By Lemma 4.3, a.s. no geodesic of \mathfrak{d}_h hits 0. In the case of a free-boundary GFF or a thick quantum wedge, it is clear that a.s. no geodesic of \mathfrak{d}_h hits ∞ . In the case of a single bead of a thin wedge, this follows since $(\mathbb{H}, h, 0, \infty) \stackrel{d}{=} (\mathbb{H}, h, \infty, 0)$. It follows that we can find $\rho = \rho(p) > 1$ such that with probability at least $1 - p$, every geodesic from z_1 to z_2 is contained in \mathbb{W}'_ρ . Let E be the event that this is the case.

By Lemma 2.2, on E the set of \mathfrak{d}_h -geodesics from z_1 to z_2 coincides with the set of $\mathfrak{d}_{h|_{\mathbb{W}'_\rho}}$ -geodesics from z_1 to z_2 . The law of the field $h|_{\mathbb{W}'_\rho}$ is absolutely continuous with respect to the law of the corresponding restriction of an appropriate embedding into $(\mathbb{H}, 0, \infty)$ of a quantum disk, which by [MS16a, Corollary 1.5] is isometric to a Brownian disk. By [BM15, Lemma 18], we infer that a.s. there is only one $\mathfrak{d}_{h|_{\mathbb{W}'_\rho}}$ -geodesic from z_1 to z_2 and that this geodesic a.s. does not hit $\partial \mathbb{D}$. Since $\mathbb{P}[E]$ can be made arbitrarily close to 1 by choosing ρ sufficiently large, we obtain the statement of the lemma. \square

In the setting of Theorem 1.3, there is no boundary to worry about but we still need to ensure that geodesics stay away from the origin, since, due to the presence of a log singularity, the restriction of the field h to a neighborhood of the origin is not absolutely continuous with respect to a quantum disk (or sphere). The following lemma addresses this issue. We do not give a proof since the argument is essentially the same as the proof of Lemma 4.1.

Lemma 4.4. *Suppose that h is either a whole-plane GFF on plus $-\alpha \log |\cdot|$ for $\alpha < Q$, an embedding into $(\mathbb{C}, 0, \infty)$ of an α -quantum cone for $\alpha < Q$, or an embedding into $(\mathbb{C}, 0, \infty)$ of a quantum sphere. Let $R > 1$ and let z_1, z_2 be sampled uniformly from $\mu_h|_{B_R(0) \setminus B_{1/R}(0)}$, normalized to be a probability measure. Almost surely, there is a unique \mathfrak{d}_h -geodesic γ_{z_1, z_2} from z_1 to z_2 , and a.s. γ_{z_1, z_2} does not hit 0 or ∞ .*

4.2 Metric gluing of two quantum wedges

In this subsection we will prove Theorem 1.1. The proof of Theorem 1.3 is similar to that of Theorem 1.1, but simpler because we only need to consider a single complementary connected component of η (note that the proof uses Lemma 4.4 in place of Lemma 4.1). So, we will only give a detailed proof of Theorem 1.1.

Throughout, we assume we are in the setting of Theorem 1.1. Recall in particular that $(\mathbb{H}, h, 0, \infty)$ is a weight- \mathfrak{w} quantum wedge (so $\mu_h(\mathbb{H}) = \infty$) if $\mathfrak{w} \geq 4/3$, or a single bead of a thin wedge with quantum area \mathfrak{a}

and left/right quantum boundary lengths \mathfrak{l}^- and \mathfrak{l}^+ (if $\mathfrak{w} < 4/3$). We assume that the embedding of h is chosen so that the quantum mass of $\mathbb{D} \cap \mathbb{H}$ is 1 (if $\mathfrak{w} \geq 4/3$) or so that the quantum mass of $\mathbb{D} \cap \mathbb{H}$ is $\mathfrak{a}/2$ (if $\mathfrak{w} < 4/3$). This is merely for convenience as the statement of the theorem is independent of the choice of embedding. We also assume that the $\text{SLE}_{8/3}(\mathfrak{w}^- - 2; \mathfrak{w}^+ - 2)$ curve η is parameterized by quantum length with respect to h .

For a connected component $U \in \mathcal{U}^- \cup \mathcal{U}^+$, let x_U (resp. y_U) be the first (resp. last) point of ∂U hit by η . Then the quantum surfaces $(U, h|_U, x_U, y_U)$ for $U \in \mathcal{U}^\pm$ are the beads of \mathcal{W}^\pm if $\mathfrak{w}^\pm < 4/3$, or all of \mathcal{W}^\pm if $\mathfrak{w}^\pm \geq 4/3$.

We now give an outline of the proof of Theorem 1.1. For most of the proof, we will truncate on a global regularity event $G_C = G_C(u, \rho)$, which we define in Lemma 4.5 just below. After establishing that G_C holds with probability close to 1 when the parameters are chosen appropriately, our main aim will be to prove the following. There is an $\alpha > 0$ such that if G_C occurs, then for small $\delta > 0$ the \mathfrak{d}_h -geodesic γ_{z_1, z_2} between two points $z_1, z_2 \in B_\rho(0)$ sampled from $\mu_h|_{B_\rho(0)}$ hits at most $O_\delta(\delta^{-1+\alpha})$ SLE segments of the form $\eta([(k-1)\delta^2, k\delta^2])$ for $k \in \mathbb{N}$. In light of Lemma 3.2, this will allow us to construct a path from z_1 to z_2 of length approximately $\mathfrak{d}_h(z_1, z_2)$ which consists of a concatenation of finitely many segments of γ_{z_1, z_2} which are contained in U for $U \in \mathcal{U}^- \cup \mathcal{U}^+$; and $O_\delta(\delta^{1-\alpha})$ paths of length at most $\delta^{1+\alpha_\delta(1)}$ which do not cross η , each of which connects two points of $\eta([(k-1)\delta^2, k\delta^2])$ for some $k \in \mathbb{N}$. This, in turn, will imply the theorem statement.

The proof of our estimate for the number of δ^2 -length SLE segments hit by a \mathfrak{d}_h -geodesic will be accomplished in two main steps. First we will show in Lemma 4.7 that on G_C , a \mathfrak{d}_h -metric ball of radius approximately δ cannot intersect too many δ^2 -length SLE segments. Then we will obtain the desired estimate in Lemma 4.8 by bounding the probability that η gets within \mathfrak{d}_h -distance δ of a given segment of the geodesic γ_{z_1, z_2} .

We are now ready to define our regularity event G_C and show that it occurs with high probability. For the statement of the following lemma, we recall the sets \mathbb{V}_ρ and \mathbb{V}'_ρ from (4.1).

Lemma 4.5. *There exists $\beta > 0$ such that the following is true. For $u \in (0, 1)$, $\rho > 2$, and $C > 1$, let $G_C = G_C(u, \rho)$ be the event that the following hold.*

1. *For each $z \in \mathbb{V}_\rho$ and each $\delta \in (0, 1]$ with $B_\delta(z; \mathfrak{d}_h) \cap \mathbb{R} = \emptyset$, we have $\mu_h(B_\delta(z; \mathfrak{d}_h)) \leq C\delta^{4-u}$.*
2. *For each $U \in \mathcal{U}^- \cup \mathcal{U}^+$ with $U \cap \mathbb{V}_\rho \neq \emptyset$, each $z \in \overline{U} \cap \mathbb{V}_\rho$, and each $\delta \in (0, 1]$, we have $\mu_h(B_\delta(z; \mathfrak{d}_{h|_U})) \geq C^{-1}\delta^{4+u}$.*
3. *For each $U \in \mathcal{U}^- \cup \mathcal{U}^+$ with $U \cap \mathbb{V}_\rho \neq \emptyset$ and each $x, y \in \partial U \cap \mathbb{V}_\rho$,*

$$\mathfrak{d}_{h|_U}(x, y) \leq C\nu_h([x, y]_{\partial U})^{1/2}(|\log \nu_{a, \ell}([x, y]_{\partial \text{BD}_{a, \ell}})| + 1)^2. \quad (4.5)$$

4. *For each $U \in \mathcal{U}^- \cup \mathcal{U}^+$ with $U \cap \mathbb{V}_\rho \neq \emptyset$, each $x, y \in \partial U \cap \mathbb{V}_\rho$, and each $\delta \in (0, 1)$,*

$$\mu_h(B_\delta([x, y]_{\partial U}; \mathfrak{d}_{h|_U})) \geq C^{-1}\delta^{2+u}\nu_h([x, y]_{\partial U}).$$

5. *For each $z \in \mathbb{V}'_\rho$ and each $\delta \in (0, 1]$, we have $B_\delta(z; \mathfrak{d}_h) \subset B_{C\delta^\beta}(z)$.*
6. *For each $t > s > 0$ such that $\eta(s) \in \mathbb{V}_{\rho/2}$ and $|t - s| \leq C^{-1}$, we have $\eta(t) \in \mathbb{V}_\rho$.*

For each $u > 0$, $\rho > 2$, and $p \in (0, 1)$ there exists $C > 1$ such that $\mathbb{P}[G_C] \geq 1 - p$.

We remark that in (4.5) it is possible to improve the exponent of the log term to $7/4 + \zeta$ for any $\zeta > 0$ (see Lemma 3.2), but the particular exponent does not matter for our purposes so for simplicity we just take it to be 2.

Proof of Lemma 4.5. Fix $u \in (0, 1)$, $\rho > 2$, and $p \in (0, 1)$. We will deduce the statement of the lemma from the results of Section 3.2 and local absolute continuity (recall that the Brownian disk is equivalent to the quantum disk). First we need to reduce ourselves to considering only finitely many quantum surfaces, rather than the surfaces parameterized by all of the complementary connected components of η (which is an infinite set of one of \mathfrak{w}^\pm is smaller than $4/3$).

Let \mathcal{F} be the σ -algebra generated by the μ_h -areas of the ordered sequences of beads of \mathcal{W}^- and \mathcal{W}^+ (so that \mathcal{F} is trivial if $\mathfrak{w}^- \wedge \mathfrak{w}^+ \geq 4/3$). For $\epsilon > 0$, let \mathcal{U}_ϵ^\pm be the set of $U \in \mathcal{U}^\pm$ such that the following is true: we have $\mu_h(U) \geq \epsilon$, and the sum of the quantum areas of the beads of \mathcal{W}^\pm which come before U is at most ϵ^{-1} . Then each \mathcal{U}_ϵ^\pm is a finite set. Furthermore, \mathcal{F} determines which elements of \mathcal{U}^\pm belong to \mathcal{U}_ϵ^\pm so the conditional law of the surfaces $(U, h|_U, x_U, y_U)$ for $U \in \mathcal{U}_\epsilon^\pm$ given \mathcal{F} is that of a collection of independent beads of a weight- \mathfrak{w}^\pm quantum wedge with given areas (or a single weight- \mathfrak{w}^\pm quantum wedge if $\mathfrak{w}^\pm \geq 4/3$).

The boundary of each $U \in \mathcal{U}^\pm$ intersects \mathbb{R} (since η is simple) and there are only finitely many $U \in \mathcal{U}^\pm$ which intersect $B_\rho(0)$ and have diameter larger than ρ^{-1} (since η is continuous and transient [MS16d]). Consequently, we can find $\epsilon = \epsilon(\rho) > 0$ such that with probability at least $1 - p/5$, it holds that each $U \in \mathcal{U}^- \cup \mathcal{U}^+$ which intersects \mathbb{V}_ρ belongs to $\mathcal{U}_\epsilon^- \cup \mathcal{U}_\epsilon^+$. Henceforth fix such an ϵ .

For $U \in \mathcal{U}^\pm$, let $\phi_U : U \rightarrow \mathbb{H}$ be the unique conformal map which takes x_U to 0 and y_U to ∞ , normalized so that the quantum measure induced by the field $h_U := h \circ \phi_U^{-1} + Q \log |(\phi_U^{-1})'|$ assigns mass 1 to $\mathbb{D} \cap \mathbb{H}$ (if $\mu_h(U) = \infty$) or mass $\mu_h(U)/2$ to $\mathbb{D} \cap \mathbb{H}$ (if $\mu_h(U) < \infty$).

Since each of the marked points x_U, y_U for $U \in \mathcal{U}^\pm$ lies in $\mathbb{R} \cup \{\infty\}$, we infer that each ϕ_U for $U \in \mathcal{U}^\pm$ a.s. maps $U \cap \mathbb{V}_{2\rho}$ to a bounded subset of \mathbb{H} lying at positive distance from 0. Since each \mathcal{U}_ϵ^\pm is a.s. finite, we can find $\tilde{\rho} > 1$ such that with probability at least $1 - p/5$,

$$\phi_U(U \cap \mathbb{V}'_{2\rho}) \subset \mathbb{V}'_{\tilde{\rho}}, \quad \forall U \in \mathcal{U}_\epsilon^- \cup \mathcal{U}_\epsilon^+.$$

Again using the finiteness of \mathcal{U}_ϵ^\pm , we can find $a > 1 \wedge \mathfrak{a}$ and $\ell > 0$ such that with probability at least $1 - p/5$,

$$\begin{aligned} \mu_h(\mathbb{V}'_{2\rho}) &\leq a/2, & \nu_h(\partial\mathbb{V}'_{2\rho} \cap \mathbb{R}) &\leq \ell/2, \\ \mu_{h_U}(\mathbb{V}'_{2\tilde{\rho}}) &\leq a/2, & \text{and } \nu_{h_U}(\partial\mathbb{V}'_{2\tilde{\rho}} \cap \mathbb{R}) &\leq \ell/2, \quad \forall U \in \mathcal{U}_\epsilon^- \cup \mathcal{U}_\epsilon^+. \end{aligned}$$

Let $(\mathbb{H}, \tilde{h}, 0, \infty)$ be a doubly marked quantum disk with area a and left/right boundary lengths each equal to ℓ , with the embedding chosen so that $\mu_{\tilde{h}}(\mathbb{D} \cap \mathbb{H}) = 1$ (if $\mathfrak{w} \geq 4/3$) or $\mathfrak{a}/2$ (if $\mathfrak{w} < 4/3$). It is easy to see from the definitions given in [DMS14, Section 4] that on the event $\{\mu_h(\mathbb{V}'_{2\rho}) \leq a/2, \nu_h(\partial\mathbb{V}'_{2\rho} \cap \mathbb{R}) \leq \ell/2\}$ the law of $h|_{\mathbb{V}'_\rho}$ is absolutely continuous with respect to the law of the corresponding restriction of \tilde{h} . Furthermore, if we condition on \mathcal{F} then on the event

$$\{\mu_{h_U}(\mathbb{V}'_{2\tilde{\rho}}) \leq a/2, \nu_{h_U}(\partial\mathbb{V}'_{2\tilde{\rho}} \cap \mathbb{R}) \leq \ell/2\},$$

the conditional law of $h_U|_{\mathbb{V}'_\rho}$ is absolutely continuous with respect to the law of the corresponding restriction of \tilde{h} .

By [MS15c, Corollary 1.5], the space \mathbb{H} equipped with the $\sqrt{8/3}$ -LQG metric induced by \tilde{h} is isometric to the Brownian disk with area a and boundary length 2ℓ . Since $\partial\mathbb{V}_r$ a.s. lies at positive quantum distance from $\partial\mathbb{V}_{2r}$ for each $r > 1$ and similarly for \mathbb{V}'_r and \mathbb{V}'_{2r} , it follows from Lemmas 3.2, 3.3, and 3.4 that we can find $C > 1$ such that conditions 1 through 4 in the statement of the lemma hold with probability at least $1 - p/2$.

By [MS16a, Theorem 1.2] and local absolute continuity, for an appropriate choice of universal $\beta > 0$ and a large enough choice of C , condition 5 holds with probability at least $1 - p/4$. By continuity and transience of η for large enough $C > 1$ (depending on ρ) condition 6 holds with probability at least $1 - p/4$. \square

Our next lemma shows that η satisfies a reverse Hölder continuity condition on the event G_C (here we recall that η is parameterized by quantum length with respect to h).

Lemma 4.6. *For each $v \in (0, 1)$, there exists $u_0 = u_0(v) \in (0, 1)$ such that for $u \in (0, u_0]$ the following is true. Let $\rho > 2$ and $C > 1$ and let $G_C = G_C(u, \rho)$ be the event of Lemma 4.5. There exists $\epsilon_0 = \epsilon_0(C, u, v)$ such that the following is true a.s. on G_C . If $0 < a < b < \infty$ such that $b - a \leq \epsilon_0$ and $\eta([a, b]) \cap \mathbb{V}_{\rho/2} \neq \emptyset$, then*

$$\text{diam}(\eta([a, b]); \mathfrak{d}_h) \geq 7(b - a)^{(1+v)/2}.$$

Proof. Throughout the proof we assume that G_C occurs. We first observe that condition 5 in Lemma 4.5 implies that there is an $\epsilon_0 = \epsilon_0(C, \rho)$ such that for $\epsilon \in (0, \epsilon_0]$ and $z \in \mathbb{V}_{\rho/2}$,

$$B_{8\epsilon^{(1+v)/2}}(z; \mathfrak{d}_h) \subset \mathbb{V}_\rho. \tag{4.6}$$

Suppose now that $0 < a < b < \infty$ such that $b - a \leq \epsilon_0$ and $\eta([a, b]) \cap \mathbb{W}_{\rho/2} \neq \emptyset$. Let $z \in \eta([a, b]) \cap \mathbb{W}_\rho$ and write $\delta := (b - a)^{(1+v)/2}$. It follows from (4.6) that $B_{8\delta}(z; \mathfrak{d}_h)$ does not intersect \mathbb{R} , and this combined with (4.6) implies there exists $U \in \mathcal{U}^-$ such that $\eta([a, b]) \subset \partial U \cap \mathbb{W}_\rho$. By condition 4 in Lemma 4.5 and since $\mathfrak{d}_h(w_1, w_2) \leq \mathfrak{d}_{h|U}(w_1, w_2)$ for each $w_1, w_2 \in \overline{U}$ (by Lemma 2.2 and Remark 2.3),

$$\mu_h(B_\delta(\eta([a, b]); \mathfrak{d}_h)) \geq \mu_h(B_\delta(\eta([a, b]); \mathfrak{d}_{h|U})) \geq C^{-1}(b - a)^{(2+u)(1+v)/2+1}.$$

By condition 1,

$$\mu_h(B_{8\delta}(z; \mathfrak{d}_h)) \leq 8^{4-u}C(b - a)^{(4-u)(1+v)/2}.$$

If u is chosen sufficiently small relative to v , then after possibly shrinking ϵ_0 we can arrange that $8^{4-u}C(b - a)^{(4-u)(1+v)/2} < C^{-1}(b - a)^{(2+u)(1+v)/2+1}$ whenever $b - a \leq \epsilon_0$, so $B_\delta(\eta([a, b]); \mathfrak{d}_h)$ cannot be contained in $B_{8\delta}(z; \mathfrak{d}_h)$. Therefore $\eta([a, b])$ cannot be contained in $B_{7\delta}(z; \mathfrak{d}_h)$. \square

We next bound the number of δ^2 -length segments of η which can intersect a \mathfrak{d}_h -metric ball on the event G_C .

Lemma 4.7. *For each $v \in (0, 1)$, there exists $u_0 = u_0(v) \in (0, 1)$ such that for $u \in (0, u_0]$, the following is true. Let $\rho > 2$ and $C > 1$ and let $G_C = G_C(u, \rho)$ be the event of Lemma 4.5. There exists $\delta_0 = \delta_0(C, u, v) > 0$ such that the following is true almost surely on G_C . Let $z \in \mathbb{W}_{\rho/2}$ and let $\delta \in (0, \delta_0]$. Then the number of $k \in \mathbb{N}$ for which $\eta([(k-1)\delta^2, k\delta^2]) \cap B_{\delta^{1+v}}(z; \mathfrak{d}_h) \neq \emptyset$ is at most δ^{-v} .*

Proof. See Figure 5 for an illustration of the argument. Throughout the proof we assume that G_C occurs. As in the proof of Lemma 4.6, if z is as in the statement of the lemma then

$$B_{3\delta^{1+v}}(z; \mathfrak{d}_h) \subset \mathbb{W}_\rho \quad (4.7)$$

for small enough δ (depending only on C and ρ). By Lemma 4.6, if $u \in (0, 1)$ is chosen sufficiently small relative to v , then for small enough $\delta > 0$ (depending only on C , u , and v) there cannot exist $k \in \mathbb{N}$ for which $\eta([(k-1)\delta^2, k\delta^2]) \subset B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. Henceforth assume that δ and u are chosen so that the above conditions are satisfied.

Let K be the set of $k \in \mathbb{N}$ for which $\eta([(k-1)\delta^2, k\delta^2]) \cap B_{\delta^{1+v}}(z; \mathfrak{d}_h) \neq \emptyset$. By our choices of δ and u , each curve segment $\eta([(k-1)\delta^2, k\delta^2])$ for $k \in K$ also intersects $\mathbb{H} \setminus B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. We will prove an upper bound for $\#K$ by comparing $\#K$ to the cardinality of another set, and then prove an upper bound for the cardinality of this set in terms of $\mu_h(B_{2\delta^{1+v}}(z; \mathfrak{d}_h))$.

Let \mathcal{V} be the set of connected components of $\mathbb{H} \setminus B_{\delta^{1+v}}(z; \mathfrak{d}_h)$. For $V \in \mathcal{V}$, let \mathcal{O}_V be the set of connected components O of $V \setminus \eta$ such that O intersects $V \setminus B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. We first argue that

$$\#K \leq 2 \sum_{V \in \mathcal{V}} \#\mathcal{O}_V + 2. \quad (4.8)$$

To see this, fix $V \in \mathcal{V}$. Let $\{[\tau_{V,j}, \bar{\tau}_{V,j}]\}_{j \in [1, J_V]_{\mathbb{Z}}}$ be the ordered sequence of time intervals for η with the property that $\eta([\tau_{V,j}, \bar{\tau}_{V,j}]) \subset V$, $\eta([\tau_{V,j}, \bar{\tau}_{V,j}]) \cap \partial V \neq \emptyset$, and $\eta([\tau_{V,j}, \bar{\tau}_{V,j}])$ intersects $V \setminus B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. Note that the number J_V of such time intervals is finite by continuity and transience of η and since $V \setminus B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$ lies at positive Euclidean distance from ∂V .

The set V has the topology of the disk (or the complement of a disk in \mathbb{H} , in the case of the unbounded component). Since η does not hit itself each curve segment $\eta([\tau_{V,j}, \bar{\tau}_{V,j}])$ is contained in a single connected component of $V \setminus \eta([0, \tau_{V,j}])$, which also has the topology of a disk (or possibly the complement of a disk in \mathbb{H} if V is the unbounded component). Since $\eta([\tau_{V,j}, \bar{\tau}_{V,j}]) \in \partial V$, the segment $\eta([\tau_{V,j}, \bar{\tau}_{V,j}])$ divides this connected component into at least two further components. Since $\eta([\tau_{V,j}, \bar{\tau}_{V,j}])$ intersects $V \setminus B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$, each such component also intersects $V \setminus B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. It follows that $J_V \leq \#\mathcal{O}_V$.

Each interval $[(k-1)\delta^2, k\delta^2]$ for $k \in K$ intersects $[\tau_{V,j}, \bar{\tau}_{V,j}]$ for some $V \in \mathcal{V}$ and some $j \in [1, J_V]_{\mathbb{Z}}$, except possibly the first and last such intervals. Furthermore, each such interval $[\tau_{V,j}, \bar{\tau}_{V,j}]$ intersects at most two intervals of the form $[(k-1)\delta^2, k\delta^2]$ for $k \in K$. Therefore, $\#K \leq 2 \sum_{V \in \mathcal{V}} J_V + 2$, whence (4.8) holds.

Now we will prove an upper bound for the right side of (4.8). For $V \in \mathcal{V}$ and $O \in \mathcal{O}_V$, we have $\partial O \cap \partial V \neq \emptyset$, since η does not hit itself. Hence we can choose $w_O \in \partial O \cap \eta$ such that $\mathfrak{d}_h(w_O, \partial B_{\delta^{1+v}}(z)) = \mathfrak{d}_h(w_O, \partial B_{2\delta^{1+v}}(z)) = \frac{1}{2}\delta^{1+v}$. We have $O \subset U_O$ for some connected component $U_O \in \mathcal{U}^- \cup \mathcal{U}^+$ of $\mathbb{H} \setminus \eta$. Let

B_O be the open $\mathfrak{d}_{h|_{U_O}}$ -ball of radius $\frac{1}{2}\delta^{1+v}$ centered at w_O . Since η does not cross B_O and $\mathfrak{d}_{h|_{U_O}} \leq \mathfrak{d}_h$, we have

$$B_O \subset B_{\frac{1}{2}\delta^{1+v}}(w_O; \mathfrak{d}_h) \subset B_{2\delta^{1+v}}(z; \mathfrak{d}_h) \setminus B_{\delta^{1+v}}(z; \mathfrak{d}_h).$$

Therefore, $B_O \subset O$ so in particular $B_O \cap B_{O'} = \emptyset$ for distinct $O, O' \in \bigcup_{V \in \mathcal{V}} \mathcal{O}_V$.

By (4.7) and condition 2 in Lemma 4.5, for each $O \in \bigcup_{V \in \mathcal{V}} \mathcal{O}_V$ we have $\mu_h(B_O) \succeq C^{-1}\delta^{(4+u)(1+v)}$, with universal implicit constant. Therefore, condition 1 in Lemma 4.5 implies that

$$C^{-1}\delta^{(4+u)(1+v)} \sum_{V \in \mathcal{V}} \#\mathcal{O}_V \preceq \mu_h(B_{2\delta^{1+v}}(z; \mathfrak{d}_h)) \preceq C\delta^{(4-u)(1+v)}$$

with universal implicit constants. By combining this with (4.8) we get

$$\#K \preceq C^2\delta^{-2u(1+v)}.$$

After possibly shrinking u and δ , we obtain the statement of the lemma. \square




Figure 5: Illustration of the proof of Lemma 4.7. The balls $B_{\delta^{1+v}}(z; \mathfrak{d}_h)$ and $B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$ are shown in light green and light blue, respectively. We first use the upper bound for the area of $B_{3\delta^{1+v}}(z; \mathfrak{d}_h)$ to argue that on G_C , no curve segment of the form $\eta([(k-1)\delta^2, k\delta^2])$ can be completely contained in $B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. So we need to estimate the number of crossings of η between $B_{\delta^{1+v}}(z; \mathfrak{d}_h)$ and $\mathbb{H} \setminus B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. This is accomplished by considering an appropriate collection of $\mathfrak{d}_{h|_{W^\pm}}$ -balls B_O (shown in pink), one in each connected component O of $\mathbb{H} \setminus (\eta \cup B_{\delta^{1+v}}(z; \mathfrak{d}_h))$ which intersects both $\partial B_{\delta^{1+v}}(z; \mathfrak{d}_h)$ and $B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. The number of such components O (8 in the figure) provides an upper bound for the number of crossings of η between $B_{\delta^{1+v}}(z; \mathfrak{d}_h)$ and $\mathbb{H} \setminus B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. The sets B_O are disjoint and each is contained in $B_{2\delta^{1+v}}(z; \mathfrak{d}_h) \setminus B_{\delta^{1+v}}(z; \mathfrak{d}_h)$. By definition of G_C , we have a lower bound for the quantum mass of each B_O and an upper bound for the quantum mass of $B_{2\delta^{1+v}}(z; \mathfrak{d}_h)$. By comparing these bounds, we obtain an upper bound on the number of sets B_O .

Next we bound the number of segments of η with quantum length δ^2 which are hit by a \mathfrak{d}_h -geodesic between two typical points.

Lemma 4.8. *Let $v \in (0, 1)$ and let $u_0 = u_0(v)$ be chosen as in Lemma 4.7. Let $u \in (0, u_0]$, $\rho > 2$, and $C > 1$ and let $G_C = G_C(u, \rho)$ be as in Lemma 4.5. Let $z_1, z_2 \in \mathbb{V}_\rho$ and let γ_{z_1, z_2} be a \mathfrak{d}_h -geodesic from z_1 to z_2 which is contained in $\mathbb{V}_{\rho/2}$, all chosen in a manner which is independent from η (viewed as a curve modulo*

parameterization). For $\delta \in (0, \delta_0]$, let $\mathcal{K}_{z_1, z_2}^\delta$ be the set of $k \in \mathbb{N}$ for which $\gamma_{z_1, z_2} \cap \eta([(k-1)\delta^2, k\delta^2]) \neq \emptyset$. There is an exponent $\alpha > 0$ such that for $\delta \in (0, 1)$, we have

$$\mathbb{E}[\#\mathcal{K}_{z_1, z_2}^\delta \mathbb{1}_{G_C} \mid h, \gamma_{z_1, z_2}] \preceq \delta^{-1-2v+\alpha(1+v)} \mathfrak{d}_h(z_1, z_2) \quad (4.9)$$

where the implicit constant in \preceq is deterministic and depends only on u, v, C , and ρ .

Proof. By possibly increasing the implicit constant in (4.9), it suffices to prove the estimate for $\delta \in (0, \delta_0]$, where $\delta_0 > 0$ depends only on u, v, C , and ρ . Fix z_1, z_2 as in the statement of the lemma. Let $N := \lfloor \delta^{-1-v} \mathfrak{d}_h(z_1, z_2) \rfloor + 1$. For $j \in [1, N-1]_{\mathbb{Z}}$ let $t_j := j\delta^{1+v}$ and let $t_N := \mathfrak{d}_h(z_1, z_2)$. Also let $V_j := B_{\delta^{1+v}}(\gamma_{z_1, z_2}(t_j); \mathfrak{d}_h)$. Since γ_{z_1, z_2} travels one unit of quantum distance in one unit of time, the union of the balls V_j covers γ_{z_1, z_2} and the intersection of any four such balls is empty. Let $\mathcal{J}_{z_1, z_2}^\delta$ be the number of $j \in [1, N]_{\mathbb{Z}}$ for which $\eta \cap V_j \neq \emptyset$. Lemma 4.7 implies that for small enough δ (depending only on u, v , and C), it holds on G_C that there are at most δ^{-v} elements of $\mathcal{K}_{z_1, z_2}^\delta$ which intersect each of the sets V_j . Hence for any such δ ,

$$\#\mathcal{K}_{z_1, z_2}^\delta \leq 3\delta^{-v} \#\mathcal{J}_{z_1, z_2}^\delta. \quad (4.10)$$

In light of (4.10), it remains only to show that

$$\mathbb{E}[\#\mathcal{J}_{z_1, z_2}^\delta \mathbb{1}_{G_C} \mid h, \gamma_{z_1, z_2}] \preceq \mathfrak{d}_h(z_1, z_2) \delta^{-1-v+\alpha(1+v)} \quad (4.11)$$

for appropriate $\alpha > 0$ as in the statement of the lemma. To this end, define the balls V_j as above. By condition 5 in Lemma 4.5 (recall that we have assumed that $\gamma_{z_1, z_2} \subset \mathbb{V}_{\rho/2}$), for each $j \in [1, N]_{\mathbb{Z}}$, the metric ball V_j is contained in the Euclidean ball \tilde{V}_j of radius $C\delta^{\beta(1+v)}$ and the same center as V_j , where $\beta > 0$ is as in condition 5. It is easy to see that there is a universal constant $\alpha_0 > 0$ such that for each $w \in \mathbb{V}_{\rho/2}$ and each $\epsilon \in (0, 1)$, we have

$$\mathbb{P}[\eta \cap B_\epsilon(w) \neq \emptyset] \preceq \epsilon^{\alpha_0},$$

with the implicit constant depending only on ρ . Indeed, this follows by stopping η at the first time it hits each of the balls $B_{\delta^{-k}}(w)$ for $k \in \mathbb{N}$ and applying the domain Markov property and transience of $\text{SLE}_\kappa(\rho^L; \rho^R)$ (the latter property is [MS16d, Theorem 1.3]).² Since η is independent from h and γ_{z_1, z_2} , we find that for each $j \in [1, N]_{\mathbb{Z}}$,

$$\mathbb{P}[\eta \cap \tilde{V}_j \neq \emptyset \mid h, \gamma_{z_1, z_2}] \preceq \delta^{\alpha_0 \beta(1+v)}.$$

By summing over all $j \in [1, N]_{\mathbb{Z}}$ we obtain (4.11) with $\alpha = \alpha_0 \beta$. \square

The following lemma shows that the metric \mathfrak{d}_h is equal to the quotient metric in Theorem 1.1 at quantum typical points.

Lemma 4.9. Fix $R > 1$. Let z_1, z_2 be sampled uniformly from $\mu_h|_{\mathbb{V}_R}$, normalized to be a probability measure. Let $\tilde{\mathfrak{d}}_h$ be the quotient metric on \mathbb{H} obtained by gluing the metric spaces $(U, \mathfrak{d}_h|_U)$ according to the natural identification. Almost surely, we have

$$\mathfrak{d}_h(z_1, z_2) = \tilde{\mathfrak{d}}_h(z_1, z_2).$$

Proof. See Figure 6 for an illustration of the proof. Fix $p \in (0, 1)$. Also let α be as in Lemma 4.5, let $v \in (0, \alpha/100)$, and let $u \in (0, u_0]$ where $u_0 = u_0(v)$ is as in Lemma 4.7. Let γ_{z_1, z_2} be the \mathfrak{d}_h -geodesic from z_1 to z_2 , which by Lemma 4.1 is a.s. unique and a.s. does not intersect $\mathbb{R} \cup \{\infty\}$. We can find $\rho > 2$ such that with probability at least $1 - p/5$,

$$\gamma_{z_1, z_2} \subset \mathbb{V}_{\rho/2} \quad \text{and} \quad \mathfrak{d}_h(z_1, z_2) \leq \rho.$$

Let E_ρ be the event that this is the case. Also choose $C > 0$ such that for this choice of ρ and u as above, the event $G_C = G_C(u, \rho)$ occurs with probability at least $1 - p/5$.

By taking the expectation of the estimate from Lemma 4.8, we obtain, in the notation of that lemma,

$$\mathbb{E}[\#\mathcal{K}_{z_1, z_2} \mathbb{1}_{G_C \cap E_\rho}] \preceq \delta^{-1-2v+\alpha(1+v)}$$

²By [Bef08, Proposition 4], $\alpha_0 = 2/3$ when $\rho^L = \rho^R = 0$, and we expect, but do not prove, that this is true for general $\rho^L, \rho^R > -2$.




Figure 6: Illustration of the proof of Lemma 4.9. We sample z_1, z_2 uniformly from $\mu_h|_{\mathbb{V}_R}$ and consider the \mathfrak{d}_h -geodesic γ_{z_1, z_2} (blue) from z_1 to z_2 , which is typically contained in $\mathbb{V}_{\rho/2}$ for some $\rho > R$. We divide the segment of η (red) which is contained in $\mathbb{V}_{\rho/2}$ into increments of quantum length δ^2 . Lemma 4.8 implies that on G_C , the number of such segments which are hit by γ_{z_1, z_2} is at most $\delta^{-1+\alpha'}$ for a small positive constant α' . On the other hand, condition 3 in the definition of G_C (recall Lemma 4.5) implies that for each δ^2 -length segment of η which is hit by γ_{z_1, z_2} , we can find a complementary connected component $U \in \mathcal{U}^-$ and a $\mathfrak{d}_{h|U}$ -geodesic between the first and last points of this segment which are hit by γ_{z_1, z_2} whose length is at most $\delta^{1+o_\delta(1)}$. These one-sided geodesics are shown in purple. We then concatenate these one-sided geodesics with the segments of γ_{z_1, z_2} between the times when it hits η to obtain a path from z_1 to z_2 which crosses η only finitely many times and whose length is at most $\mathfrak{d}_h(z_1, z_2) + o_\delta(1)$.

with the implicit constant depending only on u, v, C , and ρ . By Markov's inequality, there exists $\delta_0 = \delta_0(u, v, C, \rho) > 0$ such that for $\delta \in (0, \delta_0]$, it holds with probability at least $1 - p/2$ that $E_\rho \cap G_C$ occurs and

$$\#\mathcal{K}_{z_1, z_2}^\delta \leq \delta^{-1-3v+\alpha(1+v)} \leq \delta^{-1+\alpha/2}, \quad (4.12)$$

where the second inequality follows from our choice of v .

Now fix $\delta \in (0, \delta_0]$ and assume that $E_\rho \cap G_C$ and the event in (4.12) occur. We will bound $\tilde{\mathfrak{d}}_h(z_1, z_2)$ by constructing a path from z_1 to z_2 which is a concatenation of finitely many paths which are contained in the closure of a single set U for $U \in \mathcal{U}^- \cup \mathcal{U}^+$.

By condition 6 in Lemma 4.5, if we take $\delta \leq C^{-1/2}$ then $\eta([(k-1)\delta^2, k\delta]) \subset \mathbb{V}_\rho$ for each $k \in \mathcal{K}_{z_1, z_2}^\delta$. In particular, $\eta([(k-1)\delta^2, k\delta])$ does not intersect \mathbb{R} , so $\eta([(k-1)\delta^2, k\delta]) \subset \partial U_k$ for some $U_k \in \mathcal{U}^-$ with $U_k \cap \mathbb{V}_\rho \neq \emptyset$.

For $k \in \mathcal{K}_{z_1, z_2}^\delta$, let τ_k (resp. σ_k) be the first (resp. last) time γ_{z_1, z_2} hits $\eta([(k-1)\delta^2, k\delta])$. Let $\tilde{\gamma}_k$ be the $\mathfrak{d}_{h|U_k}$ -geodesic from $\gamma_{z_1, z_2}(\tau_k)$ to $\gamma_{z_1, z_2}(\sigma_k)$. By condition 3 in Lemma 4.5, on G_C it is a.s. the case that for this choice of U_k , we have $\text{diam}(\eta([(k-1)\delta^2, k\delta^2]); \mathfrak{d}_{h|U_k}) \leq 4C\delta(|\log \delta| + 1)^2$, so in particular

$$\text{len}(\tilde{\gamma}_k; \mathfrak{d}_{h|U_k}) \leq 4C\delta(|\log \delta| + 1)^2, \quad \forall k \in \mathcal{K}_{z_1, z_2}^\delta. \quad (4.13)$$

Let k_1 be the element of $\mathcal{K}_{z_1, z_2}^\delta$ with τ_{k_1} minimal (i.e. τ_{k_1} is the first time γ_{z_1, z_2} hits η). Inductively, if $j \in [2, \#\mathcal{K}_{z_1, z_2}^\delta]_{\mathbb{Z}}$ and k_{j-1} has been defined, let $k \in \mathcal{K}_{z_1, z_2}^\delta$ be chosen so that τ_k is the smallest time τ_k for $k \in \mathcal{K}_{z_1, z_2}^\delta$ which satisfies $\tau_k \geq \sigma_{k_{j-1}}$, if such a k exists; and otherwise let $k_j = \infty$. Let J be the smallest $j \in \mathbb{N}$ for which $k_j = \infty$. Let $\dot{\gamma}_1 := \gamma_{z_1, z_2}|_{[0, \tau_{k_1}]}$, $\dot{\gamma}_J := \gamma_{z_1, z_2}|_{[\sigma_J, \mathfrak{d}_h(z_1, z_2)]}$, and for $j \in [2, J-1]_{\mathbb{Z}}$ let $\dot{\gamma}_j := \gamma_{z_1, z_2}|_{[\sigma_{j-1}, \tau_j]}$. Then each curve $\dot{\gamma}_j$ for $j \in [1, J]_{\mathbb{Z}}$ does not hit η except at its endpoints so is contained in the closure of a single element $\tilde{U}_j \in \mathcal{U}^- \cup \mathcal{U}^+$, and its $\mathfrak{d}_{h|\tilde{U}_j}$ -length is the same as its \mathfrak{d}_h -length.

Let $\tilde{\gamma}$ be the curve obtained by concatenating the curves $\dot{\gamma}_1, \tilde{\gamma}_{k_1}, \dots, \dot{\gamma}_{k_{J-1}}, \tilde{\gamma}_J, \dot{\gamma}_J$. Then $\tilde{\gamma}$ is a path from

z_1 to z_2 and the quotient metric $\tilde{\mathfrak{d}}_h$ satisfies

$$\begin{aligned}\tilde{\mathfrak{d}}_h(z_1, z_2) &\leq \sum_{j=1}^{J-1} \text{len}(\tilde{\gamma}_{k_j}; \mathfrak{d}_{h|_{U_{k_j}}}) + \sum_{j=1}^J \text{len}(\tilde{\gamma}_j; \mathfrak{d}_{h|_{U_j}}) \\ &\leq \sum_{j=1}^{J-1} \text{len}(\tilde{\gamma}_{k_j}; \mathfrak{d}_{h|_{U_{k_j}}}) + \mathfrak{d}_h(z_1, z_2) \\ &\leq O_\delta \left(\delta^{\alpha/2} (|\log \delta| + 1)^2 \right) + \mathfrak{d}_h(z_1, z_2).\end{aligned}$$

where the last inequality is by (4.13). Sending $\delta \rightarrow 0$ shows that $\tilde{\mathfrak{d}}_h(z_1, z_2) \leq \mathfrak{d}_h(z_1, z_2)$. By Remark 2.3 we have $\tilde{\mathfrak{d}}_h(z_1, z_2) \geq \mathfrak{d}_h(z_1, z_2)$, so in fact $\tilde{\mathfrak{d}}_h(z_1, z_2) = \mathfrak{d}_h(z_1, z_2)$. Since $p \in (0, 1)$ can be made arbitrarily small, we conclude. \square

Proof of Theorem 1.1. Lemma 4.9 implies that a.s. $\mathfrak{d}_h(z_1, z_2) = \tilde{\mathfrak{d}}_h(z_1, z_2)$ for each pair (z_1, z_2) in a subset of $\mathbb{H} \times \mathbb{H}$ which is dense in $\mathbb{H} \times \mathbb{H}$ with respect to the metric $\mathfrak{d}_h \times \mathfrak{d}_h$ and dense in $U \times U$ with respect to $\mathfrak{d}_{h|_U} \times \mathfrak{d}_{h|_U}$ for each $U \in \mathcal{U}^- \cup \mathcal{U}^+$. Hence, the following is true a.s. Suppose given $\epsilon > 0$ and $w_1, w_2 \in \mathbb{H}$. Choose $U_1, U_2 \in \mathcal{U}^- \cup \mathcal{U}^+$ with $w_1 \in U_1$ and $w_2 \in U_2$. Then we can find $z_1 \in U_1$ and $z_2 \in U_2$ such that $\mathfrak{d}_h(z_1, z_2) = \tilde{\mathfrak{d}}_h(z_1, z_2)$, $\mathfrak{d}_{h|_{U_1}}(z_1, w_1) \leq \epsilon$, and $\mathfrak{d}_{h|_{U_2}}(z_2, w_2) \leq \epsilon$. Note that the latter two estimates and the triangle inequality imply that $\mathfrak{d}_h(z_1, z_2) = \tilde{\mathfrak{d}}_h(z_1, z_2) \leq \mathfrak{d}_h(w_1, w_2) + 2\epsilon$. By another application of the triangle inequality, we have $\tilde{\mathfrak{d}}_h(w_1, w_2) \leq \tilde{\mathfrak{d}}_h(z_1, z_2) + 2\epsilon \leq \mathfrak{d}_h(w_1, w_2) + 4\epsilon$, so since ϵ is arbitrary $\tilde{\mathfrak{d}}_h(w_1, w_2) \leq \mathfrak{d}_h(w_1, w_2)$ and hence $\tilde{\mathfrak{d}}_h(w_1, w_2) = \mathfrak{d}_h(w_1, w_2)$. \square

4.3 Metric gluing in the peanosphere

In this subsection we will deduce Theorem 1.6 and Theorem 1.7 from Theorem 1.1 and Theorem 1.3.

Proof of Theorem 1.6. Let η^L (resp. η^R) be the left (resp. right) boundary of $\eta'((-\infty, 0])$. Note that $\mathcal{U}^- \cup \mathcal{U}^+$ is precisely the set of connected components of $\mathbb{C} \setminus (\eta^L \cup \eta^R)$.

By [DMS14, Footnote 9] and [MS13, Theorem 1.1], η^L has the law of a chordal $\text{SLE}_{8/3}(-2/3)$ from 0 to ∞ in \mathbb{H} . By [DMS14, Theorem 1.12], the surface $(\mathbb{C} \setminus \eta^L, h|_{\mathbb{C} \setminus \eta^L}, 0, \infty)$ has the law of a weight-4/3 quantum wedge. By Theorem 1.3, it is a.s. the case that $(\mathbb{C}, \mathfrak{d}_h)$ is the metric space quotient of $(\mathbb{C} \setminus \eta^L, \mathfrak{d}_{h|_{\mathbb{C} \setminus \eta^L}})$ under the natural identification.

By [MS13, Theorem 1.11], the conditional law of η^R given η^L is that of a chordal $\text{SLE}_{8/3}(-4/3; -4/3)$ from 0 to ∞ in $\mathbb{C} \setminus \eta^L$. By Theorem 1.1, a.s. $(\mathbb{C} \setminus \eta^L, \mathfrak{d}_{h|_{\mathbb{C} \setminus \eta^L}})$ is the metric quotient of the disjoint union of the metric spaces $(\bar{U}, \mathfrak{d}_{h|_U})$ for $U \in \mathcal{U}^- \cup \mathcal{U}^+$ under the natural identification, except that we don't identify points on η^L . The theorem statement follows by combining this with the previous paragraph. \square

Next we will deduce Theorem 1.7 from Theorem 1.6 and an absolute continuity argument.

Proof of Theorem 1.7. Since η' is parameterized by μ_h and a.s. hits μ_h -a.e. point of \mathbb{C} exactly once, it follows that $\eta'(\mathfrak{t})$ is independent from η' and its conditional law given h is μ_h . Hence we can assume that (\mathbb{C}, h, ∞) is embedded into \mathbb{C} in such a way that $\eta'(\mathfrak{t}) = 0$, in which case $(\mathbb{C}, h, 0, \infty)$ is a doubly marked quantum sphere. We also assume that our embedding is such that $\mu_h(\mathbb{D}) = 1/2$.

For $R > 1$, let A_R be the closed annulus $\bar{B}_R(0) \setminus B_{1/R}(0)$. Let \mathcal{U}_R be the set of $U \in \mathcal{U}^- \cup \mathcal{U}^+$ such that $U \subset A_R$. We observe that our choice of embedding implies that U is determined by η' , viewed as a curve modulo monotone re-parameterization. Let $W_R := \bigcup_{U \in \mathcal{U}_R} U$. Almost surely, 0 is not contained in ∂U for any $U \in \mathcal{U}^- \cup \mathcal{U}^+$ and every element of $\mathcal{U}^- \cup \mathcal{U}^+$ is bounded. It follows that

$$\text{for each } R > 1, \text{ there exists } R' > 1 \text{ such that } A_R \subset W_{R'}. \quad (4.14)$$

Let $(\mathbb{C}, \hat{h}, 0, \infty)$ be a $\sqrt{8/3}$ -quantum cone independent from η' with the embedding chosen so that $\mu_{\hat{h}}(\mathbb{D}) = 1/2$ and let $\hat{\eta}'$ be the curve obtained by parameterizing $\hat{\eta}'$ by $\mu_{\hat{h}}$ -mass in such a way that $\hat{\eta}'(0) = 0$. Then for each $R > 1$, the laws of $h|_{A_R}$ and $\hat{h}|_{A_R}$ are mutually absolutely continuous, so also the laws of $(h|_{A_R}, \mathcal{U}_R)$ and $(\hat{h}|_{A_R}, \mathcal{U}_R)$ are mutually absolutely continuous.

By Theorem 1.6 and the definition of the quotient metric, it is a.s. the case that for each $\epsilon > 0$ and each $z, w \in \mathbb{C}$, there exists $N \in \mathbb{N}$, $U_1, \dots, U_N \in \mathcal{U}^- \cup \mathcal{U}^+$, and $\mathfrak{d}_{\hat{h}|_{U_i}}$ -geodesics $\hat{\gamma}_i$ for $i \in [1, N]_{\mathbb{Z}}$ whose concatenation is a path from z to w such that

$$\sum_{i=1}^N \text{len}\left(\hat{\gamma}_i; \mathfrak{d}_{\hat{h}|_{U_i}}\right) \leq \mathfrak{d}_{\hat{h}}(z, w) + \epsilon.$$

Note that if $z \in W_R$ and $\mathfrak{d}_{\hat{h}}(z, w) \leq \frac{1}{4}\mathfrak{d}_{\hat{h}}(z, \partial W_R)$, then for small enough ϵ each of the sets U_i for $i \in [1, N]_{\mathbb{Z}}$ must belong to \mathcal{U}_R .

For $R > 1$ and $z \in W_R$, let $\rho_R(z) := \mathfrak{d}_h(z, \partial W_R)$. Since metric balls are locally determined by the field, we find that for each $z \in A_R$, both $\rho_R(z)$ and the restriction of \mathfrak{d}_h to $B_{\rho_R(z)}(z; \mathfrak{d}_h)$ are a.s. determined by $(h|_{A_R}, \mathcal{U}_R)$. Consequently, it follows from the preceding paragraph and absolute continuity that a.s. for each $z, w \in W_R$ with $\mathfrak{d}_h(z, w) \leq \frac{1}{4}\rho_R(z)$ and each $\epsilon > 0$, there exists $N \in \mathbb{N}$, $U_1, \dots, U_N \in \mathcal{U}_R$, and $\mathfrak{d}_{h|_{U_i}}$ -geodesics γ_i for $i \in [1, N]_{\mathbb{Z}}$ whose concatenation is a path from z to w such that

$$\sum_{i=1}^N \text{len}\left(\gamma_i; \mathfrak{d}_{h|_{U_i}}\right) \leq \mathfrak{d}_h(z, w) + \epsilon. \quad (4.15)$$

Now let z_1, z_2 be sampled uniformly from $\mu_h|_{A_R}$, normalized to be a probability measure and let γ be the \mathfrak{d}_h -geodesic from z_1 to z_2 . By Lemma 4.4, γ is a.s. unique and a.s. does not hit 0 or ∞ . Hence (4.14) implies that for each $p \in (0, 1)$ there exists $R' > R$ such that

$$\mathbb{P}[\gamma \subset W_{R'}] \geq 1 - p.$$

On the event $\{\gamma \subset W_{R'}\}$, there a.s. exists $M \in \mathbb{N}$ and times $0 = t_0 < t_1 < \dots < t_M = \mathfrak{d}_h(z_1, z_2)$ such that $\mathfrak{d}_h(\gamma(t_i), \partial W_{R'}) \geq 4(t_i - t_{i-1})$ for each $i \in [1, M]_{\mathbb{Z}}$. By applying (4.15) for each $i \in [1, M]_{\mathbb{Z}}$ with $(z, w) = (\gamma(t_{i-1}), \gamma(t_i))$ and ϵ/M in place of ϵ , we a.s. obtain a finitely many paths, each of which is a $\mathfrak{d}_{h|_U}$ -geodesic for some $U \in \mathcal{U}_{R'}$, whose concatenation is a path from z_1 to z_2 and whose total length is at most $\mathfrak{d}_h(z_1, z_2) + \epsilon$. Since $R > 1$, $p \in (0, 1)$, and $\epsilon > 0$ are arbitrary and by (4.14), we infer that if $\tilde{\mathfrak{d}}_h$ is the quotient metric on \mathbb{C} obtained by identifying the metric spaces $(U, \mathfrak{d}_{h|_U})$ for $U \in \mathcal{U}^- \cup \mathcal{U}^+$ as in the statement of the lemma, then a.s. $\mathfrak{d}_h(z_1, z_2) = \tilde{\mathfrak{d}}_h(z_1, z_2)$. By the same argument used to conclude the proof of Theorem 1.1, we obtain the desired result. \square

References

- [BBG12] G. Borot, J. Bouttier, and E. Guitter. A recursive approach to the $O(n)$ model on random maps via nested loops. *Journal of Physics A Mathematical General*, 45(4):045002, February 2012, [1106.0153](#).
- [Bef08] V. Beffara. The dimension of the SLE curves. *Ann. Probab.*, 36(4):1421–1452, 2008, [math/0211322](#). [MR2435854 \(2009e:60026\)](#)
- [Bet15] J. Bettinelli. Scaling limit of random planar quadrangulations with a boundary. *Ann. Inst. Henri Poincaré Probab. Stat.*, 51(2):432–477, 2015, [1111.7227](#). [MR3335010](#)
- [BG09] J. Bouttier and E. Guitter. Distance statistics in quadrangulations with a boundary, or with a self-avoiding loop. *J. Phys. A*, 42(46):465208, 44, 2009, [0906.4892](#). [MR2552016](#)
- [BM15] J. Bettinelli and G. Miermont. Compact Brownian surfaces I. Brownian disks. *ArXiv e-prints*, July 2015, [1507.08776](#).
- [BMR16] E. Baur, G. Miermont, and G. Ray. Classification of scaling limits of uniform quadrangulations with a boundary. *ArXiv e-prints*, August 2016, [1608.01129](#).
- [Car15] A. Caraceni. *The geometry of large outerplanar and half-planar maps*. PhD thesis, Scuola Normale Superiore, 2015.

- [CC16] A. Caraceni and N. Curien. Self-Avoiding Walks on the UIPQ. *ArXiv e-prints*, September 2016, [1609.00245](#).
- [CL14] N. Curien and J.-F. Le Gall. The Brownian plane. *J. Theoret. Probab.*, 27(4):1249–1291, 2014, [1204.5921](#). [MR3278940](#)
- [DMS14] B. Duplantier, J. Miller, and S. Sheffield. Liouville quantum gravity as a mating of trees. *ArXiv e-prints*, September 2014, [1409.7055](#).
- [DS11] B. Duplantier and S. Sheffield. Liouville quantum gravity and KPZ. *Invent. Math.*, 185(2):333–393, 2011, [1206.0212](#). [MR2819163](#) (2012f:81251)
- [GKMW16] E. Gwynne, A. Kassel, J. Miller, and D. B. Wilson. Active spanning trees with bending energy on planar maps and SLE-decorated Liouville quantum gravity for $\kappa \geq 8$. *ArXiv e-prints*, March 2016, [1603.09722](#).
- [GM16a] E. Gwynne and J. Miller. Convergence of the self-avoiding walk on random quadrangulations to SLE $_{8/3}$ on $\sqrt{8/3}$ -Liouville quantum gravity. *ArXiv e-prints*, August 2016, [1608.00956](#).
- [GM16b] E. Gwynne and J. Miller. Scaling limit of the uniform infinite half-plane quadrangulation in the Gromov-Hausdorff-Prokhorov-uniform topology. *ArXiv e-prints*, August 2016, [1608.00954](#).
- [JS00] P. W. Jones and S. K. Smirnov. Removability theorems for Sobolev functions and quasiconformal maps. *Ark. Mat.*, 38(2):263–279, 2000. [MR1785402](#) (2001i:46049)
- [KMSW15] R. Kenyon, J. Miller, S. Sheffield, and D. B. Wilson. Bipolar orientations on planar maps and SLE $_{12}$. *ArXiv e-prints*, November 2015, [1511.04068](#).
- [Le 10] J.-F. Le Gall. Geodesics in large planar maps and in the Brownian map. *Acta Math.*, 205(2):287–360, 2010, [0804.3012](#). [MR2746349](#) (2012b:60272)
- [Le 13] J.-F. Le Gall. Uniqueness and universality of the Brownian map. *Ann. Probab.*, 41(4):2880–2960, 2013, [1105.4842](#). [MR3112934](#)
- [Le 14] J.-F. Le Gall. Random geometry on the sphere. *ArXiv e-prints*, March 2014, [1403.7943](#).
- [LSW03] G. Lawler, O. Schramm, and W. Werner. Conformal restriction: the chordal case. *J. Amer. Math. Soc.*, 16(4):917–955 (electronic), 2003, [math/0209343v2](#). [MR1992830](#) (2004g:60130)
- [Mie09] G. Miermont. Random maps and their scaling limits. In *Fractal geometry and stochastics IV*, volume 61 of *Progr. Probab.*, pages 197–224. Birkhäuser Verlag, Basel, 2009. [MR2762678](#) (2012a:60017)
- [Mie13] G. Miermont. The Brownian map is the scaling limit of uniform random plane quadrangulations. *Acta Math.*, 210(2):319–401, 2013, [1104.1606](#). [MR3070569](#)
- [MS13] J. Miller and S. Sheffield. Imaginary geometry IV: interior rays, whole-plane reversibility, and space-filling trees. *ArXiv e-prints*, February 2013, [1302.4738](#).
- [MS15a] J. Miller and S. Sheffield. An axiomatic characterization of the Brownian map. *ArXiv e-prints*, June 2015, [1506.03806](#).
- [MS15b] J. Miller and S. Sheffield. Liouville quantum gravity and the Brownian map I: The QLE(8/3,0) metric. *ArXiv e-prints*, July 2015, [1507.00719](#).
- [MS15c] J. Miller and S. Sheffield. Liouville quantum gravity spheres as matings of finite-diameter trees. *ArXiv e-prints*, June 2015, [1506.03804](#).
- [MS16a] J. Miller and S. Sheffield. Liouville quantum gravity and the Brownian map II: geodesics and continuity of the embedding. *ArXiv e-prints*, May 2016, [1605.03563](#).

- [MS16b] J. Miller and S. Sheffield. Liouville quantum gravity and the Brownian map III: the conformal structure is determined. *ArXiv e-prints*, August 2016, [1608.05391](#).
- [MS16c] J. Miller and S. Sheffield. Quantum Loewner Evolution. *Duke Mathematics Journal*, to appear, 2016, [1312.5745](#).
- [MS16d] J. Miller and S. Sheffield. Imaginary geometry I: interacting SLEs. *Probab. Theory Related Fields*, 164(3-4):553–705, 2016, [1201.1496](#). [MR3477777](#)
- [MW14] J. Miller and H. Wu. Intersections of SLE paths: the double and cut point dimension of SLE. *Probab. Theory Related Fields*, 2014, [1303.4725](#). To appear.
- [RS05] S. Rohde and O. Schramm. Basic properties of SLE. *Ann. of Math. (2)*, 161(2):883–924, 2005, [math/0106036](#). [MR2153402 \(2006f:60093\)](#)
- [RY99] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition, 1999. [MR1725357 \(2000h:60050\)](#)
- [Sch97] G. Schaeffer. Bijective census and random generation of Eulerian planar maps with prescribed vertex degrees. *Electron. J. Combin.*, 4(1):Research Paper 20, 14 pp. (electronic), 1997. [MR1465581 \(98g:05074\)](#)
- [Sch00] O. Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118:221–288, 2000, [math/9904022](#). [MR1776084 \(2001m:60227\)](#)
- [Ser97] L. Serlet. A large deviation principle for the Brownian snake. *Stochastic Process. Appl.*, 67(1):101–115, 1997. [MR1445046](#)
- [She07] S. Sheffield. Gaussian free fields for mathematicians. *Probab. Theory Related Fields*, 139(3-4):521–541, 2007. [MR2322706 \(2008d:60120\)](#)
- [She16a] S. Sheffield. Conformal weldings of random surfaces: SLE and the quantum gravity zipper. *Annals of Probability*, to appear, 2016, [1012.4797](#).
- [She16b] S. Sheffield. Quantum gravity and inventory accumulation. *Annals of Probability*, to appear, 2016, [1108.2241](#).
- [SS13] O. Schramm and S. Sheffield. A contour line of the continuum Gaussian free field. *Probab. Theory Related Fields*, 157(1-2):47–80, 2013, [math/0605337](#). [MR3101840](#)
- [SW05] O. Schramm and D. B. Wilson. SLE coordinate changes. *New York J. Math.*, 11:659–669 (electronic), 2005, [math/0505368](#). [MR2188260 \(2007e:82019\)](#)
- [Ver12] R. Vershynin. Introduction to the non-asymptotic analysis of random matrices. In *Compressed sensing*, pages 210–268. Cambridge Univ. Press, Cambridge, 2012. [MR2963170](#)